arXiv:0909.5250v1 [math.DG] 29 Sep 2009

Genericity of Caustics and Wavefronts on an r-corner

Takaharu Tsukada[1]

College of Humanities & Sciences, Department of Mathematics,
Nihon University

Abstract

We investigate genericities of reticular Lagrangian maps and reticular Legendrian maps in order to give generic classifications of caustics and wavefronts generated by a hypersurface germ without or with a boundary in a smooth manifold. We also give simpler proofs of main results in [9],[10].

1 Introduction

Lagrangian and Legendrian singularities can be found in many problems of differential geometry, calculus of variations and mathematical physics. One of the most successful their applications are the study of singularities of caustics and wavefronts. For example, the particles incident along geodesics from a smooth hypersurface in a Riemannian manifold to conormal directions define a Lagrangian submanifold at a point in the cotangent bundle and define a Legendrian submanifolds at a point in the 1-jet bundle. The caustic generated by the hypersurface is regarded as the caustic of the Lagrangian map defined by the restriction of the cotangent bundle projection to the Lagrangian submanifold and the wavefront generated by the hypersurface is regarded as the wavefront of the Legendrian map defined by the restriction of the 1-jet bundle projection to the Legendrian submanifold. Therefore the studies of the caustics and wavefronts generated by smooth hypersurfaces are reduced to the studies of Lagrangian and Legendrian singularities.

In [9] and [10] we investigated the more general cases when the hypersurface has a boundary, a corner, or *an r-corner*. In these cases particles incident from each edge of the hypersurface gives a *symplectic regular r-cubic configuration* at a point of the cotangent bundle which is a generalisation of the notion of Lagrangian submanifolds and particles incident from each edge of the hypersurface gives a *contact regular r-cubic configuration* at a point of the 1-jet bundle which is a generalisation of the notion of Legendrian submanifolds. The caustic generated by the hypersurface germ with an r-corner is given by the caustic of the symplectic regular r-cubic configuration which is a generalisation of the notion of *quasi-caustics* given by S.Janeszko (cf., [6]). In these papers we investigated the stabilities of caustics and wavefronts generated by the hypersurface with an r-corner by studying the stabilities of *reticular Lagrangian, Legendrian maps* which are generalisations of the notions of Lagrangian, Legendrian maps for our situations.

In this paper, we investigate the genericities of caustics and wavefronts generated by a hypersurface with an r-corner. In order to realize this purpose, we shall investigate the genericities of reticular Lagrangian, Legendrian maps. In these processes, we shall need to prove that the *stabilities* and the *transversally stabilities* of reticular Lagrangian, Legendrian maps are equivalent respectively. These proofs are simpler than that of the assertions (1)⇔(5) of Theorem 5.5 in [9, p.587] and Theorem 7.4 in [10, p.123] respectively.

---

[1] Higashijujo 3-1-16 Kita-ku, Tokyo 114-0001 JAPAN. e-mail : tsukada@math.chs.nihon-u.ac.jp

The main results in this paper are generic classification of caustics generated by the hypersurface with an r-corner on an n dimensional manifold in the cases $n \leq 5, r = 0$ and $n \leq 3, r = 1$ and a generic classification of wavefronts generated by the hypersurface with an r-corner on an n dimensional manifold in the cases $n \leq 6, r = 0$ and $n \leq 4, r = 1$. In order to realize this, we shall classify generic reticular Lagrangian, Legendrian maps for the above cases.

By our theory, we have that: A generic caustics is one of the types $A_2, A_3^{\pm}, A_4, A_5^{\pm}, A_6, D_4^{\pm}, D_4^{\pm}, D_6^{\pm}, E_6^{\pm}$ in the case $n \leq 5, r = 0$ and $B_2^{\pm}, B_3^{\pm}, B_4^{\pm}, C_3^{\pm}, C_4^{\pm}, F_4^{\pm}$ in the case $n \leq 3, r = 1$. A generic wavefront is one of the types $A_1, A_2, A_3, A_4, A_4, A_6, D_4^{\pm}, D_5, D_6^{\pm}, E_6$ in the case $n \leq 6, r = 0$ and $B_2, B_3, B_4, C_3^{\pm}, C_4, F_4$ in the case $n \leq 4, r = 1$.



Figure 1: the caustic F_4^+ and the wavefront C_3^+

This paper consists of three parts. In Part I, we recall stabilities of unfoldings under the equivalence relations *reticular* $\mathcal{P}$-$\mathcal{R}^+$*-equivalence* and *reticular* $\mathcal{P}$-$\mathcal{K}$*-equivalence*. They work as the equivalence relations of generating families of reticular Lagrangian, Legendrian maps respectively. In part II and III we recall that the equivalence relations of reticular Lagrangian, Legendrian maps are reduced to that of their generating families. In part IV we shall study that the genericities of reticular Lagrangian, Legendrian maps are reduced to that of their generating families.

Part I

Stability of unfoldings

2 Preliminaries

Let $\mathbb{H}^r = \{(x_1, \ldots, x_r) \in \mathbb{R}^r | x_1 \geq 0, \ldots, x_r \geq 0\}$ be an r-corner. We denote by $\mathcal{E}(r; k_1, r; k_2)$ the set of all germs at 0 in $\mathbb{H}^r \times \mathbb{R}^{k_1}$ of smooth maps $\mathbb{H}^r \times \mathbb{R}^{k_1} \to \mathbb{H}^r \times \mathbb{R}^{k_2}$ and set $\mathfrak{M}(r; k_1, r; k_2) = \{f \in \mathcal{E}(r; k_1, r; k_2) | f(0) = 0\}$. We denote $\mathcal{E}(r; k_1, k_2)$ for $\mathcal{E}(r; k_1, 0; k_2)$ and denote $\mathfrak{M}(r; k_1, k_2)$ for $\mathfrak{M}(r; k_1, 0; k_2)$.

If $k_2 = 1$ we write simply $\mathcal{E}(r;k)$ for $\mathcal{E}(r;k,1)$ and $\mathfrak{M}(r;k)$ for $\mathfrak{M}(r;k,1)$. We also write $\mathcal{E}(k)$ for $\mathcal{E}(0;k)$ and $\mathfrak{M}(k)$ for $\mathfrak{M}(0;k)$. Then $\mathcal{E}(r;k)$ is an $\mathbb{R}$-algebra in the usual way and $\mathfrak{M}(r;k)$ is its unique maximal ideal.

Let $I_r = \{1,2,\dots,r\}$ and $(x,y) = (x_1,\dots,x_r,y_1,\dots,y_k)$ be a fixed coordinate system of $(\mathbb{H}^r \times \mathbb{R}^k, 0)$. We denote by $\mathcal{B}(r;k)$ the group of diffeomorphism germs on $(\mathbb{H}^r \times \mathbb{R}^k, 0)$ of the form:

$$\phi(x,y) = (x_1\phi_1^1(x,y),\dots,x_r\phi_1^r(x,y),\phi_2^1(x,y),\dots,\phi_2^k(x,y)).$$

We also denote by $\mathcal{B}_n(r;k+n)$ the group of diffeomorphism germs on $(\mathbb{H}^r \times \mathbb{R}^{k+n}, 0)$ of the form:

$$\phi(x,y,u) = (x_1\phi_1^1(x,y,u),\dots,x_r\phi_1^r(x,y,u),\phi_2^1(x,y,u),\dots,\phi_2^k(x,y,u),\phi_3^1(u),\dots,\phi_3^n(u)).$$

We denote $J^l(r+k,1)$ the set of l-jets at 0 of germs in $\mathfrak{M}(r;k)$ and let $\pi_l : \mathfrak{M}(r;k) \to J^l(r+k,1)$ be the natural projection. We denote $j^l f(0)$ the l-jet of $f \in \mathfrak{M}(r;k)$. We also denote $\phi(x,y,u) = (x\phi_1(x,y,u),\phi_2(x,y,u),\phi_3(u))$ and denote other notations analogously. In this paper all maps and all map germs are supposed to be smoothly.

3 Reticular $\mathcal{P}$-$\mathcal{R}^+$-stability of unfoldings

We recall the stabilities of unfoldings under *the reticular $\mathcal{R}^+$-equivalence* which is developed in [9]. In order to distinguish equivalence relations between function germs in $\mathfrak{M}(r;k)$ and their unfoldings, we denote this equivalence relation by *the reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalence* in this paper.

We say that $f, g \in \mathcal{E}(r;k)$ are *reticular $\mathcal{R}$-equivalent* if there exists $\phi \in \mathcal{B}(r;k)$ such that $g = f \circ \phi$.

We say that function germs $f(x,y_1,\dots,y_{k_1}) \in \mathfrak{M}(r;k_1)^2$ and $g(x,y_1,\dots,y_{k_2}) \in \mathfrak{M}(r;k_2)^2$ are *stably reticular $\mathcal{R}$-equivalent* if f and g are reticular $\mathcal{R}$-equivalent after additions of non-degenerate quadratic forms in the variables y.

We say that a function germ $f \in \mathfrak{M}(r;k)$ is *reticular $\mathcal{R}$-l-determined* if all function germ which has the same l-jet of f is reticular $\mathcal{R}$-equivalent to f.

Lemma 3.1 (cf., [9, Lemma 4.2]) *Let $f(x,y) \in \mathfrak{M}(r;k)$ and let*

$$\mathfrak{M}(r;k)^{l+1} \subset \mathfrak{M}(r;k)(\langle x_1\frac{\partial f}{\partial x_1},\dots,x_r\frac{\partial f}{\partial x_r}\rangle + \mathfrak{M}(r;k)\langle\frac{\partial f}{\partial y_1},\dots,\frac{\partial f}{\partial y_k}\rangle) + \mathfrak{M}(r;k)^{l+2},$$

then f is reticular $\mathcal{R}$-l-determined. Conversely if $f(x,y) \in \mathfrak{M}(r;k)$ be reticular $\mathcal{R}$-l-determined, then

$$\mathfrak{M}(r;k)^{l+1} \subset \langle x_1\frac{\partial f}{\partial x_1},\dots,x_r\frac{\partial f}{\partial x_r}\rangle_{\mathcal{E}(r;k)} + \mathfrak{M}(r;k)\langle\frac{\partial f}{\partial y_1},\dots,\frac{\partial f}{\partial y_k}\rangle.$$

We denote $x\frac{\partial f}{\partial x}$ for $(x_1\frac{\partial f}{\partial x_1},\dots,x_r)\frac{\partial f}{\partial x_r}$ and $\frac{\partial f}{\partial y}$ for $(\frac{\partial f}{\partial y_1},\dots,\frac{\partial f}{\partial y_k})$, and denote other notations analogously.

Let $F(x,y,u) \in \mathfrak{M}(r;k+n_1)$, $G(x,u,v) \in \mathfrak{M}(r;k+n_2)$ be unfoldings of $f \in \mathfrak{M}(r;k)$. We say that G *is reticular $\mathcal{P}$-$\mathcal{R}^+$-f-induced from F* if there exist $\Phi \in \mathfrak{M}(r;k+n_2,r;k+n_1)$ and $\alpha \in \mathfrak{M}(n_2)$ satisfying the following conditions:
(1) $\Phi(x,y,0) = (x,y,0)$ for all $(x,y) \in (\mathbb{H}^r \times \mathbb{R}^k, 0)$,

(2) Φ can be written in the form: $\Phi(x,y,v)=(x\phi_1(x,y,v),\phi_2(x,y,v),\phi_3(v))$,
(3) $G(x,y,v)=F\circ\Phi(x,y,v)+\alpha(v)$ for all $(x,y,v)\in(\mathbb{H}^r\times\mathbb{R}^{k+n_2},0)$.

We say that $F,G\in\mathcal{E}(r;k+n)$ are *reticular $\mathcal{P}$-$\mathcal{R}^{(+)}$-equivalent* if there exist $\Phi\in\mathcal{B}_n(r;k+n)$ (and $\alpha\in\mathcal{E}(n)$) such that $G=F\circ\Phi(+\alpha)$. We call (Φ,α) a reticular $\mathcal{P}$-$\mathcal{R}^{(+)}$-isomorphism from F to G.

Definition 3.2 Here we recall the definitions of several stabilities of unfoldings under the reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalence. Let $F(x,y,u)\in\mathfrak{M}(r;k+n)$ be an unfolding of $f(x,y)\in\mathfrak{M}(r;k)$.

We say that F is *reticular $\mathcal{P}$-$\mathcal{R}^+$-stable* if the following condition holds: For any neighbourhood U of 0 in $\mathbb{R}^{r+k+n}$ and any representative $\tilde{F}\in C^\infty(U,\mathbb{R})$ of F, there exists a neighbourhood $N_{\tilde{F}}$ of $\tilde{F}$ in C^∞-topology such that for any element $\tilde{G}\in N_{\tilde{F}}$ the germ $\tilde{G}|_{\mathbb{H}^r\times\mathbb{R}^{k+n}}$ at $(0,y_0,u_0)$ is reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent to F for some $(0,y_0,u_0)\in U$.

We say that F is *reticular $\mathcal{P}$-$\mathcal{R}^+$-versal* if all unfolding of f is reticular $\mathcal{P}$-$\mathcal{R}^+$-f-induced from F.

We say that F is *reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimally versal* if

$$\mathcal{E}(r;k)=\langle x\frac{\partial f}{\partial x},\frac{\partial f}{\partial y}\rangle_{\mathcal{E}(r;k)}+\langle 1,\frac{\partial F}{\partial u}|_{u=0}\rangle_{\mathbb{R}}.$$

We say that F is *reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimally stable* if

$$\mathcal{E}(r;k+n)=\langle x\frac{\partial F}{\partial x},\frac{\partial F}{\partial y}\rangle_{\mathcal{E}(r;k+n)}+\langle 1,\frac{\partial F}{\partial u}\rangle_{\mathcal{E}(n)}.$$

We say that F is *reticular $\mathcal{P}$-$\mathcal{R}^+$-homotopically stable* if for any smooth path-germ $(\mathbb{R},0)\to\mathcal{E}(r;k+n),t\mapsto F_t$ with $F_0=F$, there exists a smooth path-germ $(\mathbb{R},0)\to\mathcal{B}_n(r;k+n)\times\mathcal{E}(n),t\mapsto(\Phi_t,\alpha_t)$ with $(\Phi_0,\alpha_0)=(id,0)$ such that each (Φ_t,α_t) is a reticular $\mathcal{P}$-$\mathcal{R}^+$-isomorphism from F to F_t, that is $F_t=F\circ\Phi_t+\alpha_t$ for t around 0.

Theorem 3.3 (cf., [9, Theorem 4.5]) *Let $F\in\mathfrak{M}(r;k+n)$ be an unfolding of $f\in\mathfrak{M}(r;k)$. Then the following are equivalent.*
(1) *F is reticular $\mathcal{P}$-$\mathcal{R}^+$-stable.*
(2) *F is reticular $\mathcal{P}$-$\mathcal{R}^+$-versal.*
(3) *F is reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimally versal.*
(4) *F is reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimally stable.*
(5) *F is reticular $\mathcal{P}$-$\mathcal{R}^+$-homotopically stable.*

For a function germ $f(x,y)\in\mathfrak{M}(r;k)$, if $1,a_1,\ldots,a_n\in\mathcal{E}(r;k)$ is a representative of a basis of the vector space

$$\mathcal{E}(r;k)/\langle x\frac{\partial f}{\partial x},\frac{\partial f}{\partial y}\rangle_{\mathcal{E}(r;k)},$$

then the function germ $f+a_1u_1+\cdots+a_nu_n\in\mathfrak{M}(r;k+n)$ is a reticular $\mathcal{P}$-$\mathcal{R}^+$-stable unfolding of f. We call n the reticular $\mathcal{R}^+$-codimension of f.

We call a function germ $f \in \mathfrak{M}(r;k)$ is *$\mathcal{R}$-simple* if the following holds: For a sufficiently higher integer l, there exists a neighbourhood N of $j^l f(0)$ in $J^l(r+k,1)$ such that N intersects finite $\mathcal{R}$-orbits. By §17.4[1, p.279] we have that:

Theorem 3.4 *An $\mathcal{R}$-simple function germ in $\mathfrak{M}(1;k)^2$ is stably $\mathcal{R}$-equivalent to one of the following function germ:*

$$B_l^{\pm} : \pm x^l \ (l \geq 2), \quad C_l^{\pm} : xy \pm y^l \ (l \geq 3), \quad F_4^{\pm} : \pm x^2 + y^3.$$

4 Reticular $\mathcal{P}$-$\mathcal{K}$-stability of unfoldings

We recall the stabilities of unfoldings under *the reticular $\mathcal{K}$-equivalence* which is developed in [10]. In this paper we denote this equivalence relation by *the reticular $\mathcal{P}$-$\mathcal{K}$-equivalence.*

We say that $f, g \in \mathcal{E}(r;k)$ are *reticular $\mathcal{K}$-equivalent* if there exist $\phi \in \mathcal{B}(r;k)$ and a unit $a \in \mathcal{E}(r;k)$ such that $g = a \cdot f \circ \phi$.

We say that function germs $f(x, y_1, \ldots, y_{k_1}) \in \mathfrak{M}(r;k_1)^2$ and $g(x, y_1, \ldots, y_{k_2}) \in \mathfrak{M}(r;k_2)^2$ are *stably reticular $\mathcal{K}$-equivalent* if f and g are reticular $\mathcal{K}$-equivalent after additions of non-degenerate quadratic forms in the variables y.

We say that a function germ $f \in \mathfrak{M}(r;k)$ is *reticular $\mathcal{K}$-l-determined* if all function germ which has the same l-jet of f is reticular $\mathcal{K}$-equivalent to f.

Lemma 4.1 (cf., [10, Lemma 6.2]) *Let $f(x,y) \in \mathfrak{M}(r;k)$ and let*

$$\mathfrak{M}(r;k)^{l+1} \subset \mathfrak{M}(r;k)(\langle f, x\frac{\partial f}{\partial x}\rangle + \mathfrak{M}(r;k)\langle \frac{\partial f}{\partial y}\rangle) + \mathfrak{M}(r;k)^{l+2},$$

then f is reticular $\mathcal{K}$-l-determined. Conversely if $f(x,y) \in \mathfrak{M}(r;k)$ be reticular $\mathcal{K}$-l- determined, then

$$\mathfrak{M}(r;k)^{l+1} \subset \langle f, x\frac{\partial f}{\partial x}\rangle_{\mathcal{E}(r;k)} + \mathfrak{M}(r;k)\langle \frac{\partial f}{\partial y}\rangle.$$

Let $F(x,y,u) \in \mathfrak{M}(r;k+n_1)$, $G(x,y,v) \in \mathfrak{M}(r;k+n_2)$ be unfoldings of $f(x,y) \in \mathfrak{M}(r;k)$. We say that G *is reticular $\mathcal{P}$-$\mathcal{K}$-f-induced from F* if there exist $\Phi \in \mathfrak{M}(r;k+n_2, r;k+n_1)$ and $\alpha \in \mathcal{E}(r;k+n_2)$ satisfying the following conditions:
(1) $\Phi(x,y,0) = (x,y,0)$ and $\alpha(x,y,0) = 1$ for all $(x,y) \in (\mathbb{H}^r \times \mathbb{R}^k, 0)$,
(2) Φ can be written in the form: $\Phi(x,y,v) = (x\phi_1(x,y,v), \phi_2(x,y,v), \phi_3(v))$,
(3) $G(x,y,v) = \alpha(x,y,v) \cdot F \circ \Phi(x,y,v)$ for all $(x,y,v) \in (\mathbb{H}^r \times \mathbb{R}^{k+n_2}, 0)$

We say that $F, G \in \mathcal{E}(r;k+n)$ are *reticular $\mathcal{P}$-$\mathcal{K}$-equivalent* if there exist $\Phi \in \mathcal{B}_n(r;k+n)$ and a unit $\alpha \in \mathcal{E}(r;k+n)$ such that $G = \alpha \cdot F \circ \Phi$. We call (Φ, α) a reticular $\mathcal{P}$-$\mathcal{K}$-isomorphism from F to G.

Definition 4.2 Here we recall the definitions of several stabilities of unfoldings under the reticular $\mathcal{P}$-$\mathcal{K}$-equivalence. Let $F(x,y,u) \in \mathfrak{M}(r;k+n)$ be an unfolding of $f(x,y) \in \mathfrak{M}(r;k)$.

We say that F is *reticular $\mathcal{P}$-$\mathcal{K}$-stable* if the following condition holds: For any neighbourhood U of 0 in $\mathbb{R}^{r+k+n}$ and any representative $\tilde{F} \in C^\infty(U, \mathbb{R})$ of F, there exists a neighbourhood $N_{\tilde{F}}$ of $\tilde{F}$ in C^∞-topology such that for any element $\tilde{G} \in N_{\tilde{F}}$ the germ $\tilde{G}|_{\mathbb{H}^r \times \mathbb{R}^{k+n}}$ at $(0, y_0, u_0)$ is reticular $\mathcal{P}$-$\mathcal{K}$-equivalent to F for some $(0, y_0, u_0) \in U$.

We say that F is *reticular $\mathcal{P}$-$\mathcal{K}$-versal* if all unfolding of f is reticular $\mathcal{P}$-$\mathcal{K}$-f-induced from F.

We say that F is *reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimally versal* if

$$\mathcal{E}(r;k) = \langle f, x\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}\rangle_{\mathcal{E}(r;k)} + \langle \frac{\partial F}{\partial u}|_{u=0}\rangle_{\mathbb{R}}.$$

We say that F is *reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimally stable* if

$$\mathcal{E}(r;k+n) = \langle F, x\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}\rangle_{\mathcal{E}(r;k+n)} + \langle \frac{\partial F}{\partial u}\rangle_{\mathcal{E}(n)}.$$

We say that F is *reticular $\mathcal{P}$-$\mathcal{K}$-homotopically stable* if for any smooth path-germ $(\mathbb{R},0) \to \mathcal{E}(r;k+n), t \mapsto F_t$ with $F_0 = F$, there exists a smooth path-germ $(\mathbb{R},0) \to \mathcal{B}_n(r;k+n) \times \mathcal{E}(r;k+n), t \mapsto (\Phi_t, \alpha_t)$ with $(\Phi_0, \alpha_0) = (id, 1)$ such that each (Φ_t, α_t) is a reticular $\mathcal{P}$-$\mathcal{K}$-isomorphism from F to F_t, that is $F_t = \alpha_t \cdot F \circ \Phi_t$ for t around 0.

Theorem 4.3 (cf., [10, Theorem 6.5]) *Let $F \in \mathfrak{M}(r;k+n)$ be an unfolding of $f \in \mathfrak{M}(r;k)$. Then the following are equivalent.*
(1) *F is reticular $\mathcal{P}$-$\mathcal{K}$-stable.*
(2) *F is reticular $\mathcal{P}$-$\mathcal{K}$-versal.*
(3) *F is reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimally versal.*
(4) *F is reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimally stable.*
(5) *F is reticular $\mathcal{P}$-$\mathcal{K}$-homotopically stable.*

For a function germ $f(x,y) \in \mathfrak{M}(r;k)$, if $a_1, \ldots, a_n \in \mathcal{E}(r;k)$ is a representative of a basis of the vector space

$$\mathcal{E}(r;k)/\langle f, x\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}\rangle_{\mathcal{E}(r;k)},$$

then the function germ $f + a_1u_1 + \cdots + a_nu_n \in \mathfrak{M}(r;k+n)$ is a reticular $\mathcal{P}$-$\mathcal{K}$-stable unfolding of f. We call n the reticular $\mathcal{K}$-codimension of f.

We call a function germ $f \in \mathfrak{M}(r;k)$ is *$\mathcal{K}$-simple* if the following holds: For a sufficiently higher integer l, there exists a neighbourhood N of $j^l f(0)$ in $J^l(r+k,1)$ such that N intersects finite $\mathcal{K}$-orbits.

Theorem 4.4 *A $\mathcal{K}$-simple function germ in $\mathfrak{M}(1;k)^2$ is stably $\mathcal{K}$-equivalent to one of the following function germ:*

$$B_l : x^l \ (l \geq 2), \quad C_l^{\varepsilon} : xy + \varepsilon y^l \ (\varepsilon^{l-1} = 1, l \geq 3), \quad F_4 : x^2 + y^3.$$

Part II

Reticular Lagrangian maps

5 Symplectic regular r-cubic configurations

Let $(q,p) = (q_1, \ldots, q_n, p_1, \ldots, p_n)$ be a canonical coordinate system of $(T^*\mathbb{R}^n, 0)$ and $\pi : (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be the cotangent bundle equipped with the canonical symplectic structure $dp \wedge dq$. We define

$$L^0_\sigma = \{(q,p) \in (T^*\mathbb{R}^n, 0) | q_\sigma = p_{I_r - \sigma} = q_{r+1} = \cdots = q_n = 0, q_{I_r - \sigma} \geq 0\}$$

for $\sigma \subset I_r$.

Definition 5.1 Let $\{L_\sigma\}_{\sigma \subset I_r}$ be a family of 2^r Lagrangian submanifold germs on $(T^*\mathbb{R}^n, 0)$. Then $\{L_\sigma\}_{\sigma \subset I_r}$ is called *a symplectic regular r-cubic configuration* if there exists a symplectic diffeomorphism germ S on $(T^*\mathbb{R}^n, 0)$ such that $L_\sigma = S(L^0_\sigma)$ for all $\sigma \subset I_r$.

6 Reticular Lagrangian maps and their generating families

We introduce a main result in [9] about the relations of reticular Lagrangian maps and their generating families.

Let $\mathbb{L} = \{(q,p) \in T^*\mathbb{R}^n | q_1 p_1 = \cdots = q_r p_r = q_{r+1} = \cdots = q_n = 0, q_{I_r} \geq 0\}$ be the representative as a germ of the union of L^0_σ for $\sigma \subset I_r$. We call a map germ

$$(\mathbb{L}, 0) \xrightarrow{i} (T^*\mathbb{R}^n, 0) \xrightarrow{\pi} (\mathbb{R}^n, 0)$$

a reticular Lagrangian map if there exists a symplectic diffeomorphism germ S on $(T^*\mathbb{R}^n, 0)$ such that $i = S|_{\mathbb{L}}$. We call i *the reticular Lagrangian embedding* of $\pi \circ i$. We call S *an extension* of i and call $\{i(L^0_\sigma)\}_{\sigma \subset I_r}$ *the symplectic regular r-cubic configuration* associated with $\pi \circ i$.

Caustics: Let $\pi \circ i$ be a reticular Lagrangian map. Let C_σ be the caustic of the Lagrangian map $\pi \circ i|_{L^0_\sigma}$ for $\sigma \subset I_r$, that is, the set of critical values of $\pi \circ i|_{L^0_\sigma}$. Let $Q_{\sigma,\tau} = \pi \circ i(L^0_\sigma \cap L^0_\tau)$ for $\sigma \neq \tau \subset I_r$. We define the *caustic* of $\pi \circ i$ by

$$\bigcup_{\sigma \subset I_r} C_\sigma \cup \bigcup_{\sigma \neq \tau} Q_{\sigma,\tau}.$$

We remark that for $\tau_1, \tau_2 \subset I_r$ $(\tau_1 \neq \tau_2)$ we have $Q_{\tau_1,\tau_2} \subset Q_{\sigma, \sigma \cup \{i\}}$, where $\sigma = \tau_1 \cap \tau_2$ and i be any element of $(\tau_1 - \sigma) \cup (\tau_2 - \sigma)$. This means that $\bigcup_{\sigma \neq \tau} Q_{\sigma,\tau}$ is equal to the union of $Q_{\sigma,\tau}$ for $\sigma \subset \tau \subset I_r$, $\#(\tau - \sigma) = 1$. For example, in the case $r = 2$ we have

$$\bigcup_{\sigma \neq \tau} Q_{\sigma,\tau} \ = \ Q_{\emptyset,1} \cup Q_{\emptyset,2} \cup Q_{1,\{1,2\}} \cup Q_{2,\{1,2\}}.$$

A function germ $F(x,y,q) \in \mathfrak{M}^2(r; k+n)$ is called *S-non-degenerate* if

$$x_1, \ldots, x_r, \frac{\partial F}{\partial x_1}, \ldots, \frac{\partial F}{\partial x_r}, \frac{\partial F}{\partial y_1}, \ldots, \frac{\partial F}{\partial y_k}$$

are independent on $(\mathbb{H}^k \times \mathbb{R}^{k+n}, 0)$, that is

$$\mathrm{rank} \begin{pmatrix} \dfrac{\partial^2 F}{\partial x \partial y} & \dfrac{\partial^2 F}{\partial x \partial u} \\ \dfrac{\partial^2 F}{\partial y \partial y} & \dfrac{\partial^2 F}{\partial y \partial u} \end{pmatrix}_0 = r + k.$$

Definition 6.1 Let $\{L_\sigma\}_{\sigma\subset I_r}$ be a symplectic regular r-cubic configuration in $(T^*\mathbb{R}^n,0)$ and $F(x,y,q)\in\mathfrak{M}(r;k+n)^2$ be a function germ. We call F *a generating family* of $\{L_\sigma\}_{\sigma\subset I_r}$ if the following conditions hold:
(1) F is S-non-degenerate,
(2) $F|_{x_\sigma=0}$ is a generating family of L_σ for $\sigma\subset I_r$, that is

$$L_\sigma=\{(q,\frac{\partial F}{\partial q}(x,y,q))\in(T^*\mathbb{R}^n,0)|x_\sigma=\frac{\partial F}{\partial x_{I_r-\sigma}}=\frac{\partial F}{\partial y}=0,x_{I_r-\sigma}\geq 0\}.$$

We also call F a generating family of a reticular Lagrangian map $\pi\circ i$ if F is a generating family of the symplectic regular r-cubic configuration $\{i(L^0_\sigma)\}_{\sigma\subset I_r}$.

We call a symplectic diffeomorphism germ ϕ on $(T^*\mathbb{R}^n,0)$ *a reticular diffeomorphism* if $\phi(L^0_\sigma)=L^0_\sigma$ for $\sigma\subset I_r$. We say that reticular Lagrangian maps $\pi\circ i_1,\pi\circ i_2:(\mathbb{L},0)\to(T^*\mathbb{R}^n,0)\to(\mathbb{R}^n,0)$ are *Lagrangian equivalent* if there exist a reticular diffeomorphism ϕ and a Lagrangian equivalence Θ of π such that the following diagram is commutative:

$$\begin{tikzcd}
(\mathbb{L},0) \arrow[r, "i_1"] \arrow[d, "\phi|_{\mathbb{L}}"'] & (T^*\mathbb{R}^n,0) \arrow[r, "\pi"] \arrow[d, "\Theta"'] & (\mathbb{R}^n,0) \arrow[d, "g"'] \\
(\mathbb{L},0) \arrow[r, "i_2"] & (T^*\mathbb{R}^n,0) \arrow[r, "\pi"] & (\mathbb{R}^n,0),
\end{tikzcd}$$

where g is the diffeomorphism of the base space of π induced from Θ.

We remark that there is not the condition that a reticular diffeomorphism is a symplectic diffeomorphism in the definition in [9]. But a reticular diffeomorphism defined in [9] consists of a restriction to $\mathbb{L}$ of compositions of two symplectic diffeomorphism and a Lagrangian equivalence, it follows that a reticular diffeomorphism in [9] is automatically a restriction to $\mathbb{L}$ of symplectic diffeomorphism. So our definition is equivalent to that in [9].

We say that function germs $F(x,y_1,\ldots,y_{k_1},q)\in\mathfrak{M}(r;k_1+n)$ and $G(x,y_1,\ldots,y_{k_2},q)\in\mathfrak{M}(r;k_2+n)$ are *stably reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent* if F and G are reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent after additions of non-degenerate quadratic forms in the variables y.

Theorem 6.2 (cf., [9, Theorem 3.2]) (1) *For any reticular Lagrangian map $\pi\circ i$, there exists a function germ $F\in\mathfrak{M}(r;k+n)^2$ which is a generating family of $\pi\circ i$.*
(2) *For any S-non-degenerate function germ $F\in\mathfrak{M}(r;k+n)^2$, there exists a reticular Lagrangian map of which F is a generating family.*
(3) *Two reticular Lagrangian maps are Lagrangian equivalent if and only if their generating families are stably reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent.*

7 Stabilities of reticular Lagrangian maps

We introduce several stabilities of reticular Lagrangian maps:
Stability: For any open set U in $T^*\mathbb{R}^n$, we denote $S(U,T^*\mathbb{R}^n)$ the space of symplectic embeddings from U to $T^*\mathbb{R}^n$ with C^∞-topology. We say that a reticular Lagrangian map $\pi\circ i:(\mathbb{L}^0,0)\to(T^*\mathbb{R}^n,0)\to(\mathbb{R}^n,0)$ is *stable* if the following holds: For any extension S of i and any representative $\tilde{S}\in S(U,T^*\mathbb{R}^n)$ of S, there exists a neighbourhood $N_{\tilde{S}}$ of $\tilde{S}$ such that for any $\tilde{S}'\in N_{\tilde{S}}$ the reticular Lagrangian maps $\pi\circ\tilde{S}'_{w_0}|_{\mathbb{L}^0}$ and $\pi\circ i$ are Lagrangian equivalent for some $w_0=(0,\ldots,0,p^0_{r+1},\ldots,p^0_n)\in U$, where the symplectic diffeomorphism

germ $\tilde{S}'_{w_0}$ on $(T^*\mathbb{R}^n, 0)$ is defined by $w \mapsto \tilde{S}'(w + w_0) - \tilde{S}'(w_0)$.

Homotopically stability: Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a reticular Lagrangian map. A one-parameter family of symplectic diffeomorphisms $\bar{S} : (T^*\mathbb{R}^n \times \mathbb{R}, (0,0)) \to (T^*\mathbb{R}^n, 0)((Q, P, t) \mapsto S_t(Q, P))$ is called a *reticular Lagrangian deformation* of i if $S_0|_{\mathbb{L}} = i_0$. Let ϕ be a reticular diffeomorphism on $(T^*\mathbb{R}^n, 0)$. A map germ $\bar{\phi} : (T^*\mathbb{R}^n \times \mathbb{R}, (0,0)) \to (T^*\mathbb{R}^n, 0)((Q, P, t) \mapsto \phi_t(Q, P))$ is called a *one-parameter deformation of reticular diffeomorphisms* of ϕ if $\phi_0 = \phi$ and ϕ_t is a reticular diffeomorphism for t around 0. We say that a reticular Lagrangian map $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ is *homotopically stable* if for any reticular Lagrangian deformation $\bar{S} = \{S_t\}$ of i , there exists a one-parameter deformation of reticular diffeomorphisms $\bar{\phi} = \{\phi_t\}$ of $id_{T^*\mathbb{R}^n}$ and a one-parameter family of Lagrangian equivalences $\bar{\Theta} = \{\Theta_t\}$ of π with $\Theta_0 = id_{T^*\mathbb{R}^n}$ such that $S_t = \Theta_t \circ S_0 \circ \phi_t$ for t around 0.

Infinitesimally stability: Let S be a symplectic diffeomorphism on $(T^*\mathbb{R}^n, 0)$. We say that a vector field v on $(T^*\mathbb{R}^n, 0)$ along S is *an infinitesimal symplectic transformation of* S if there exists a reticular Lagrangian deformation $\bar{S} = \{S_t\}$ such that $S_0 = S$ and $\frac{dS_t}{dt}|_{t=0} = v$. We say that a vector field ξ on $(T^*\mathbb{R}^n, 0)$ is *an infinitesimally reticular diffeomorphism* if there exists a one-parameter deformation of reticular diffeomorphisms $\bar{\phi} = \{\phi_t\}$ of $id_{T^*\mathbb{R}^n}$ such that $\frac{d\phi_t}{dt}|_{t=0} = \xi$. We say that a vector field η on $(T^*\mathbb{R}^n, 0)$ is *an infinitesimally Lagrangian equivalence* if there exists a one-parameter family of Lagrangian equivalences $\bar{\Theta} = \{\Theta_t\}$ of π such that $\Theta_0 = id_{T^*\mathbb{R}^n}$ and $\frac{d\Theta_t}{dt}|_{t=0} = \eta$. We say that a reticular Lagrangian map $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ is *infinitesimally stable* if for any extension S of i and any infinitesimally symplectic transformation v of S, there exists an infinitesimally reticular diffeomorphism ξ and an infinitesimally Lagrange equivalence η such that $v = S_* \xi + \eta \circ S$.

We say that a function germ H on $(T^*\mathbb{R}^n, 0)$ is *fiber preserving* if H has the form $H(q, p) = \sum_{j=1}^n h_j(q) p_j + h_0(q)$.

We recall the following theorem which is proved in [9].

Theorem 7.1 (cf., [9, p.587 Theorem 5.5]) *Let* $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ *be a reticular Lagrangian map with a generating family* $F(x, y, q) \in \mathfrak{M}(r; k+n)^2$. *Then the following are equivalent.*
(u) F *is a reticular* $\mathcal{R}^+$*-stable unfolding of* $F|_{q=0}$.
(hs) $\pi \circ i$ *is homotopically stable.*
(is) $\pi \circ i$ *is infinitesimally stable.*
(a) *For any function germ* f *on* $(T^*\mathbb{R}^n, 0)$, *there exists a fiber preserving function germ* H *on* $(T^*\mathbb{R}^n, 0)$ *such that* $f \circ i = H \circ i$.
(s) $\pi \circ i$ *is stable.*

The definition of the infinitesimally stability of reticular Lagrangian maps in [9] seem to be different from our one. But these are equivalent by Lemma 12.1. The assertion that (u), (hs),(is) and (a) are all equivalent is proved in [9]. But the proof of (u)$\Leftrightarrow$(s) is slightly complicated. So we shall prove this assertion by another way in Section 12.

Part III

Reticular Legendrian maps

8 Contact regular r-cubic configurations

Let $J^1(\mathbb{R}^n,\mathbb{R})$ be the 1-jet bundle of functions in n-variables which may be considered as $\mathbb{R}^{2n+1}$ with a natural coordinate system $(q,z,p)=(q_1,\dots,q_n,z,p_1,\dots,p_n)$, where q is a coordinate system of $\mathbb{R}^n$. We equip the contact structure on $J^1(\mathbb{R}^n,\mathbb{R})$ defined by the canonical 1-form $\theta=dz-\sum_{j=1}^n p_jdq_j$. We have a natural projection $\tilde{\pi}:J^1(\mathbb{R}^n,\mathbb{R})\to\mathbb{R}^n\times\mathbb{R}$ by $\tilde{\pi}(q,z,p)=(q,z)$.

Let $\tilde{L}^0_\sigma=\{(q,z,p)\in(J^1(\mathbb{R}^n,\mathbb{R}),0)|q_\sigma=p_{I_r-\sigma}=q_{r+1}=\cdots=q_n=z=0,q_{I_r-\sigma}\geq 0\}$ for each $\sigma\subset I_r$.

Definition 8.1 Let $\{L_\sigma\}_{\sigma\subset I_r}$ be a family of 2^r Legendrian submanifold germs on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$. Then $\{L_\sigma\}_{\sigma\subset I_r}$ is called *a contact regular r-cubic configuration* if there exists a contact diffeomorphism germ C on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ such that $L_\sigma=C(\tilde{L}^0_\sigma)$ for all $\sigma\subset I_r$.

9 Reticular Legendrian maps and their generating families

We introduce a main result in [10] about the relations of reticular Legendrian maps and their generating families.

Let $\tilde{\mathbb{L}}=\{(q,z,p)\in J^1(\mathbb{R}^n,\mathbb{R})|q_1p_1=\cdots=q_rp_r=q_{r+1}=\cdots=q_n=z=0,q_{I_r}\geq 0\}$ be a representative as a germ of the union of $\tilde{L}^0_\sigma$ for all $\sigma\subset I_r$. We call a map germ

$$(\tilde{\mathbb{L}},0)\xrightarrow{i}(J^1(\mathbb{R}^n,\mathbb{R}),0)\xrightarrow{\tilde{\pi}}(\mathbb{R}^n\times\mathbb{R},0)$$

a reticular Legendrian map if there exists a contact diffeomorphism germ C on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ such that $i=C|_{\tilde{\mathbb{L}}}$. We call i *the reticular Legendrian embedding of* $\tilde{\pi}\circ i$.

Wavefronts: Let $\tilde{\pi}\circ i$ be a reticular Legendrian map. We define the *wavefront* of $\tilde{\pi}\circ i$ by the union of $\tilde{\pi}\circ i(L^0_\sigma)$ for all $\sigma\subset I_r$.

A function germ $F(x_1,\dots,x_r,y_1,\dots,y_k,q_1,\dots,q_n,z)\in\mathfrak{M}(r;k+n+1)$ is called *C-non-degenerate* if $\frac{\partial F}{\partial x}(0)=\frac{\partial F}{\partial y}(0)=0$ and

$$x_1,\dots,x_r,F,\frac{\partial F}{\partial x_1},\dots,\frac{\partial F}{\partial x_r},\frac{\partial F}{\partial y_1},\dots,\frac{\partial F}{\partial y_k}$$

are independent on $(\mathbb{H}^r\times\mathbb{R}^{k+n+1},0)$, that is

$$\mathrm{rank}\begin{pmatrix}\frac{\partial F}{\partial y} & \frac{\partial F}{\partial q} & \frac{\partial F}{\partial z}\\ \frac{\partial^2 F}{\partial x\partial y} & \frac{\partial^2 F}{\partial x\partial q} & \frac{\partial^2 F}{\partial x\partial z}\\ \frac{\partial^2 F}{\partial y\partial y} & \frac{\partial^2 F}{\partial y\partial q} & \frac{\partial^2 F}{\partial y\partial z}\end{pmatrix}_0=r+k+1.$$

Definition 9.1 Let $\{L_\sigma\}_{\sigma\subset I_r}$ be a contact regular r-cubic configuration in $(J^1(\mathbb{R}^n,\mathbb{R}),0)$. Then a function germ $F(x,y,q,z)\in\mathfrak{M}(r;k+n+1)$ is called *a generating family* of $\{L_\sigma\}_{\sigma\subset I_r}$ if the following conditions hold:
(1) F is C-non-degenerate,
(2) For each $\sigma\subset I_r$, the function germ $F|_{x_\sigma=0}$ is a generating family of L_σ, that is

$$L_\sigma=\{(q,z,\frac{\partial F}{\partial q}/(-\frac{\partial F}{\partial z}))\in(J^1(\mathbb{R}^n,\mathbb{R}),0)|\ x_\sigma=\frac{\partial F}{\partial x_{I_r-\sigma}}=\frac{\partial F}{\partial y}=F=0,x_{I_r-\sigma}\geq 0\}.$$

We also call a function germ F a generating family of a reticular Legendrian map $\tilde{\pi}\circ i$ if F is a generating family of the contact regular r-cubic configuration $\{i(\tilde{L}^0_\sigma)\}_{\sigma\subset I_r}$.

We call a contact diffeomorphism germ ϕ on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ *a reticular diffeomorphism* if $\phi(\tilde{L}^0_\sigma)=\tilde{L}^0_\sigma$ for $\sigma\subset I_r$. We say that reticular Legendrian maps $\tilde{\pi}\circ i_1,\tilde{\pi}\circ i_2:(\tilde{\mathbb{L}},0)\to(J^1(\mathbb{R}^n,\mathbb{R}),0)\to(\mathbb{R}^n\times\mathbb{R},0)$ are *Legendrian equivalent* if there exist a reticular diffeomorphism ϕ and a Legendrian equivalence Θ of $\tilde{\pi}$ such that the following diagram is commutative:

$$\begin{tikzcd}
(\tilde{\mathbb{L}},0) \arrow[r, "i_1"] \arrow[d, "\phi|_{\tilde{\mathbb{L}}}"'] & (J^1(\mathbb{R}^n,\mathbb{R}),0) \arrow[r, "\tilde{\pi}"] \arrow[d, "\Theta"'] & (\mathbb{R}^n\times\mathbb{R},0) \arrow[d, "g"'] \\
(\tilde{\mathbb{L}},0) \arrow[r, "i_2"] & (J^1(\mathbb{R}^n,\mathbb{R}),0) \arrow[r, "\tilde{\pi}"] & (\mathbb{R}^n\times\mathbb{R},0),
\end{tikzcd}$$

where g is the diffeomorphism of the base space of $\tilde{\pi}$ induced from Θ.

As after the definition of a reticular diffeomorphism in the previous part, our definition of a reticular diffeomorphism and that in [10] are equivalent.

We say that function germs $F(x,y_1,\ldots,y_{k_1},q,z)\in\mathfrak{M}(r;k_1+n+1)$ and $F(x,y_1,\ldots,y_{k_2},q,z)\in\mathfrak{M}(r;k_2+n+1)$ are *stably reticular $\mathcal{P}$-$\mathcal{K}$-equivalent* if F and G are reticular $\mathcal{P}$-$\mathcal{K}$-equivalent after additions of non-degenerate quadratic forms in the variables y.

Theorem 9.2 (cf., [10, Theorem 5.6]) (1) *For any reticular Legendrian map $\tilde{\pi}\circ i$, there exists a function germ $F\in\mathfrak{M}(r;k+n+1)$ which is a generating family of $\tilde{\pi}\circ i$.*
(2) *For any C-non-degenerate function germ $F\in\mathfrak{M}(r;k+n+1)$, there exists a reticular Legendrian map of which F is a generating family.*
(3) *Two reticular Legendrian maps are Legendrian equivalent if and only if their generating families are stably reticular $\mathcal{P}$-$\mathcal{K}$-equivalent.*

10 Stabilities of reticular Legendrian maps

We introduce several stabilities of reticular Legendrian maps:
Stability: For any open set U in $J^1(\mathbb{R}^n,\mathbb{R})$, we denote $C(U,J^1(\mathbb{R}^n,\mathbb{R}))$ the space of contact embeddings from U to $J^1(\mathbb{R}^n,\mathbb{R})$ with C^∞-topology. We say that a reticular Legendrian map $\tilde{\pi}\circ i:(\tilde{\mathbb{L}},0)\to(J^1(\mathbb{R}^n,\mathbb{R}),0)\to(\mathbb{R}^n\times\mathbb{R},0)$ is *stable* if the following holds: For any extension C of i and any representative $\tilde{C}\in C(U,J^1(\mathbb{R}^n,\mathbb{R}))$ of C, there exists a neighbourhood $N_{\tilde{C}}$ of $\tilde{C}$ such that for any $\tilde{C}'\in N_{\tilde{C}}$ the reticular Legendrian maps $\tilde{\pi}\circ\tilde{C}'_{w_0}|_{\tilde{\mathbb{L}}}$ and $\tilde{\pi}\circ i$ are Legendrian equivalent for some $w_0=(0,\ldots,0,p^0_{r+1},\ldots,p^0_n)\in U$, where the contact diffeomorphism germ $\tilde{C}'_{w_0}$ on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ is chosen that the reticular Legendrian maps $\tilde{\pi}\circ\tilde{C}'|_{\tilde{\mathbb{L}}}:(\tilde{\mathbb{L}},w_0)\to(J^1(\mathbb{R}^n,\mathbb{R}),\tilde{C}'(w_0))\to(\mathbb{R}^n\times\mathbb{R},\tilde{\pi}\circ\tilde{C}'(w_0))$ and $\tilde{\pi}\circ\tilde{C}'_{w_0}|_{\tilde{\mathbb{L}}}:(\tilde{\mathbb{L}},0)\to(J^1(\mathbb{R}^n,\mathbb{R}),0)\to(\mathbb{R}^n\times\mathbb{R},0)$ are Legendrian equivalent.

Homotopically stability: Let $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ be a reticular Legendrian map. A one-parameter family of contact diffeomorphism germs $\bar{C} : (J^1(\mathbb{R}^n, \mathbb{R}) \times \mathbb{R}, (0, 0)) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0)((Q, Z, P, t) \mapsto C_t(Q, Z, P))$ is called a *reticular Legendrian deformation* of i if $C_0|_{\tilde{\mathbb{L}}} = i$. Let ϕ be a reticular diffeomorphism on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$. A map germ $\bar{\phi} : (J^1(\mathbb{R}^n, \mathbb{R}) \times \mathbb{R}, (0, 0)) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0)((Q, Z, P, t) \mapsto \phi_t(Q, Z, P))$ is called a *one-parameter deformation of reticular diffeomorphisms* of ϕ if $\phi_0 = \phi$ and ϕ_t is a reticular diffeomorphism for t around 0. We say that a reticular Legendrian map $\tilde{\pi} \circ i : (\mathbb{L}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ is *homotopically stable* if for any reticular Legendrian deformation $\bar{C} = \{C_t\}$ of i, there exists a one-parameter deformation of reticular diffeomorphisms $\bar{\phi} = \{\phi_t\}$ of $id_{J^1(\mathbb{R}^n, \mathbb{R})}$ such that $C_t = \Theta_t \circ C_0 \circ \phi_t$ for t around 0.

Infinitesimally stability: Let C be a contact diffeomorphism germ on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$. We say that a vector field v on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ along C is *infinitesimal contact transformation* if there exists a reticular Legendrian deformation $\bar{C} = \{C_t\}$ on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ such that $C_0 = C$ and $\frac{dC_t}{dt}|_{t=0} = v$. We say that a vector field ξ on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ is *infinitesimally reticular diffeomorphism* if there exists a one-parameter deformation of reticular diffeomorphisms $\bar{\phi} = \{\phi_t\}$ of $id_{J^1(\mathbb{R}^n, \mathbb{R})}$ such that $\frac{d\phi_t}{dt}|_{t=0} = \xi$. We say that a vector field η on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ is *infinitesimally Legendrian equivalence* if there exists a one-parameter family of Legendrian equivalences $\bar{\Theta} = \{\Theta_t\}$ on $\tilde{\pi}$ such that $\Theta_0 = id_{J^1(\mathbb{R}^n, \mathbb{R})}$ and $\frac{d\Theta_t}{dt}|_{t=0} = \eta$. We say that a reticular Legendrian map $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ is *infinitesimally stable* if for any extension C of i and any infinitesimally contact transformation v of C, there exists an infinitesimally reticular diffeomorphism ξ and an infinitesimally Legendrian equivalence η such that $v = C_* \xi + \eta \circ C$.

We say that a function germ H on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ is *fiber preserving* if H has the form $H(q, z, p) = \sum_{j=1}^n h_j(q, z) p_j + h_0(q, z)$.

Theorem 10.1 (cf., [10, p.123 Theorem 7.4]) *Let* $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ *be a reticular Legendrian map with a generating family* $F(x, y, q, z) \in \mathfrak{M}(r; k + n + 1)$. *Then the following are equivalent.*
(u) F *is a reticular* $\mathcal{P}$-$\mathcal{K}$*-stable unfolding of* $F|_{q=z=0}$.
(hs) $\tilde{\pi} \circ i$ *is homotopically stable.*
(is) $\tilde{\pi} \circ i$ *is infinitesimally stable.*
(a) *For any function germ* f *on* $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$*, there exists a fiber preserving function germ* H *on* $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ *such that* $f \circ i = H \circ i$.
(s) $\tilde{\pi} \circ i$ *is stable.*

We shall prove the assertion (u)⇔(s) by a simpler way than that of [10] in Section 13.

Part IV

Genericities of reticular Lagrangian, Legendrian maps

11 Finitely determinacy of reticular Lagrangian, Legendrian maps

Definition 11.1 Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a reticular Lagrangian map and l be a non-negative number. We say that $\pi \circ i$ is l-determined if the following condition holds: For any extension S of i, the reticular Lagrangian map $\pi \circ S'|_{\mathbb{L}}$ and $\pi \circ i$ are Lagrangian equivalent for any symplectic diffeomorphism germ S' on $(T^*\mathbb{R}^n, 0)$ satisfying $j^l S(0) = j^l S'(0)$.

Lemma 11.2 *Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a reticular Lagrangian map. Let S_1, S_2 are symplectic diffeomorphism germ on $(T^*\mathbb{R}^n, 0)$ such that $S_1|_{\mathbb{L}} = S_2|_{\mathbb{L}} = i$. Then there exists a reticular diffeomorphism germ ϕ such that $S_1 = S_2 \circ \phi$.*

Proof. Set $\phi = (S_2)^{-1} \circ S_1$. Then we have that $\phi|_{\mathbb{L}} = id$ and $S_1 = S_2 \circ \phi$ ■

By this lemma we have that the finitely determinacy of reticular Lagrangian maps do not depend on choices of extensions of their reticular Lagrangian embeddings.

Theorem 11.3 *Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a reticular Lagrangian map. If $\pi \circ i$ is infinitesimally stable then $\pi \circ i$ is $(n+1)$-determined.*

Proof. Let S be an extension of i. Since the infinitesimally stability of reticular Lagrangian maps is invariant under Lagrangian equivalences, we may assume that the canonical relation P_S associated with S has the form:

$$P_S = \{(Q, -\frac{\partial H}{\partial Q}(Q,p), -\frac{\partial H}{\partial p}(Q,p), p) \in (T^*\mathbb{R}^n \times T^*\mathbb{R}^n, (0,0))\}$$

for some function germ $H(Q,p) \in \mathfrak{M}(2n)^2$. Then $F(x,y,q) = H_0(x,y) + \langle y, q\rangle \in \mathfrak{M}(r; n+n)^2$ is a generating family of $\pi \circ i$, where $H_0(x,y) = H(x_1, \ldots, x_r, 0, \ldots, 0, y_1, \ldots, y_n) \in \mathfrak{M}(r;n)^2$. Since $\pi \circ i$ is infinitesimally stable, it follows that F is a reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimally versal unfolding of $H_0(x,y)$ by Theorem 7.1. This means that $\mathcal{E}(r;n)/\langle x\frac{\partial H_0}{\partial x}, \frac{\partial H_0}{\partial y}\rangle_{\mathcal{E}(r;n)}$ is at most $(n+1)$-dimension. It follows that

$$\mathfrak{M}(r;n)^{n+1} \subset \langle x\frac{\partial H_0}{\partial x}, \frac{\partial H_0}{\partial y}\rangle_{\mathcal{E}(r;n)}.$$

Therefore we have that

$$\mathfrak{M}(r;n)^{n+3} \subset \mathfrak{M}(r;n)(\langle x\frac{\partial H_0}{\partial x}\rangle + \mathfrak{M}(r;n)\langle \frac{\partial H_0}{\partial y}\rangle).$$

This means that H_0 is reticular $\mathcal{R}$-$(n+2)$-determined by Lemma 3.1. Let a symplectic diffeomorphism germ S' on $(T^*\mathbb{R}^n, 0)$ satisfying $j^{n+1}S(0) = j^{n+1}S'(0)$ be given. Since

$\frac{\partial p\circ S}{\partial p}(0) = \frac{\partial p\circ S'}{\partial p}(0)$, it follows that there exists a function germ $H'(Q,p) \in \mathfrak{M}(2n)^2$ such that

$$P_{S'} = \{(Q, -\frac{\partial H'}{\partial Q}(Q,p), -\frac{\partial H'}{\partial p}(Q,p), p)\}.$$

Then the function germ $G(x,y,q) := H'_0(x,y) + \langle y, q\rangle \in \mathfrak{M}(r; n+n)^2$ is a generating family of $\pi \circ S'|_{\mathbb{L}}$, where $H'_0(x,y) = H'(x,0,y) \in \mathfrak{M}(r;n)^2$. Since $j^{n+1}S(0) = j^{n+1}S'(0)$, it means that $j^{n+1}\frac{\partial H}{\partial Q}(0) = j^{n+1}\frac{\partial H'}{\partial Q}(0)$, $j^{n+1}\frac{\partial H}{\partial p}(0) = j^{n+1}\frac{\partial H'}{\partial p}(0)$. Therefore we have that $j^{n+2}H(0) = j^{n+2}H'(0)$ and hence $j^{n+2}H_0(0) = j^{n+2}H'_0(0)$. Thus there exists $\Phi(x,y) \in \mathcal{B}(r;n)$ such that $H_0 = H'_0 \circ \Phi$. We set $G'(x,y,q) := G(\Phi(x,y), q) \in \mathfrak{M}(r; n+n)^2$. Then G and G' are reticular $\mathcal{P}$-$\mathcal{R}$-equivalent, and F and G' are reticular $\mathcal{P}$-$\mathcal{R}^+$-infinitesimal versal unfoldings of $H_0(x,y)$. It follows that F and G are reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent. Therefore $\pi\circ i$ and $\pi \circ S'|_{\mathbb{L}}$ are Lagrangian equivalent. ■

We consider contact diffeomorphism germs on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$. Let (Q,Z,P) be canonical coordinates on the source space and (q,z,p) be canonical coordinates of the target space. We define the following notations:
$\imath : (J^1(\mathbb{R}^n,\mathbb{R}) \cap \{Z=0\}, 0) \to (J^1(\mathbb{R}^n,\mathbb{R}),0)$ be the inclusion map on the source space.

$$\begin{aligned}
C(J^1(\mathbb{R}^n,\mathbb{R}),0) &= \{C : (J^1(\mathbb{R}^n,\mathbb{R}),0) \to (J^1(\mathbb{R}^n,\mathbb{R}),0) | C : \text{ contact diffeomorphism}\} \\
C^\theta(J^1(\mathbb{R}^n,\mathbb{R}),0) &= \{C \in C(J^1(\mathbb{R}^n,\mathbb{R}),0) |\ C \text{ preserves the canonical 1-form } \} \\
C_Z(J^1(\mathbb{R}^n,\mathbb{R}),0) &= \{C \circ \imath\ | C \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)\} \\
C^\theta_Z(J^1(\mathbb{R}^n,\mathbb{R}),0) &= \{C \circ \imath\ | C \in C^\theta(J^1(\mathbb{R}^n,\mathbb{R}),0)\}.
\end{aligned}$$

Definition 11.4 Let $\tilde\pi\circ i : (\tilde{\mathbb{L}},0) \to (J^1(\mathbb{R}^n,\mathbb{R}),0) \to (\mathbb{R}^n\times\mathbb{R},0)$ be a reticular Legendrian map. We say that $\tilde\pi\circ i$ is l-determined if the following condition holds: For any extension $C \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ of i, the reticular Legendrian map $\tilde\pi\circ C'|_{\tilde{\mathbb{L}}}$ and $\tilde\pi\circ i$ are Legendrian equivalent for $C' \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^lC(0) = j^lC'(0)$.

Lemma 11.5 (cf., [10, p.116 Lemma 7.2]) *Let $\{L_\sigma\}_{\sigma\subset I_r}$ be a contact regular r-cubic configuration in $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ defined by $C \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$. Then there exists $C' \in C^\theta(J^1(\mathbb{R}^n,\mathbb{R}),0)$ that also defines $\{L_\sigma\}_{\sigma\subset I_r}$.*

By Lemma 11.5 we may consider the following other definition of finitely determinacy of reticular Legendrian maps:
(1) The definition given by replacing $C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ to $C^\theta(J^1(\mathbb{R}^n,\mathbb{R}),0)$.
(2) The definition given by replacing $C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ to $C_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$.
(3) The definition given by replacing $C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ to $C^\theta_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$.
Then the following holds:

Proposition 11.6 *Let $\tilde\pi\circ i : (\tilde{\mathbb{L}},0) \to (J^1(\mathbb{R}^n,\mathbb{R}),0) \to (\mathbb{R}^n\times\mathbb{R},0)$ be a reticular Legendrian map. Then*
(A) If $\tilde\pi\circ i$ is l-determined of the original definition, then $\tilde\pi\circ i$ is l-determined of the definition (1).
(B) If $\tilde\pi\circ i$ is l-determined of the definition (1), then $\tilde\pi\circ i$ is l-determined of the definition (3).

(C) *If $\tilde{\pi}\circ i$ is l-determined of the definition* (3)*, then $\tilde{\pi}\circ i$ is $(l+1)$-determined of the definition* (2)*.*
(D) *If $\tilde{\pi}\circ i$ is l-determined of the definition* (2)*, then $\tilde{\pi}\circ i$ is l-determined of the original definition.*

Proof. (A) Let $C \in C^{\theta}(J^1(\mathbb{R}^n,\mathbb{R}),0)$ be an extension of i. Let $C' \in C^{\theta}(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^lC(0) = j^lC'(0)$ be given. Since $C, C' \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$, we have that $\tilde{\pi}\circ C'|_{\tilde{\mathbb{L}}}$ and $\tilde{\pi}\circ i$ are Legendrian equivalent.
(B) Let $C \in C^{\theta}_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ be an extension of i. Let $C' \in C^{\theta}_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^lC(0) = j^lC'(0)$ be given. We define $C_1, C_1' \in C^{\theta}(J^1(\mathbb{R}^n,\mathbb{R}),0)$ by $C_1(Q,Z,P) := (q_C(Q,P), Z + z_C(Q,P), p_C(Q,P)), C_1'(Q,Z,P) := (q_{C'}(Q,P), Z + z_{C'}(Q,P), p_{C'}(Q,P))$. Then C_1 is an extension of i and we have that $j^lC_1(0) = j^lC_1'(0)$. Therefore we have that $\tilde{\pi}\circ C_1'|_{\tilde{\mathbb{L}}} = \tilde{\pi}\circ C'|_{\tilde{\mathbb{L}}}$ and $\tilde{\pi}\circ i$ are Legendrian equivalent.
(D) Let $C \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ be an extension of i. Let $C' \in C(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^lC(0) = j^lC'(0)$ be given. We set $C_1 := C|_{Z=0}, C_1' := C'|_{Z=0} \in C_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ and we have that $j^lC_1(0) = j^lC_1'(0)$. Therefore $\tilde{\pi}\circ C_1'|_{\tilde{\mathbb{L}}} = \tilde{\pi}\circ C'|_{\tilde{\mathbb{L}}}$ and $\tilde{\pi}\circ i$ are Legendrian equivalent.
(C) Let $C \in C_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ be an extension of i. Let $C' \in C_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^{l+1}C(0) = j^{l+1}C'(0)$ be given. Then there exist function germs $f(Q,P), g(Q,P) \in \mathcal{E}(2n)$ such that $C^*(dz - pdq) = -fPdQ, C'^*(dz - pdq) = -gPdQ$. Indeed f is defined by that $fP_j = -\frac{\partial z_C}{\partial Q_j} + p_C\frac{\partial q_C}{\partial Q_j}$ for $j = 1,\dots,n$. We define the diffeomorphism germs ϕ, ψ on $(J^1(\mathbb{R}^n,\mathbb{R}) \cap \{Z = 0\}, 0)$ by $\phi(Q,P) = (Q, fP), \psi(Q,P) = (Q, gP)$. We set $C_1 := C\circ\phi^{-1}, C_1' := C'\circ\psi^{-1} \in C^{\theta}_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ Then $j^l\phi(0)$ and $j^l\psi(0)$ depend only on $j^{l+1}C(0)$, therefore we have that $j^lC_1(0) = j^lC_1'(0)$. Since $\tilde{\pi}\circ i$ and $\tilde{\pi}\circ C_1|_{\tilde{\mathbb{L}}}$ are Legendrian equivalent, it follows that $\tilde{\pi}\circ C_1|_{\tilde{\mathbb{L}}}$ and $\tilde{\pi}\circ C_1'|_{\tilde{\mathbb{L}}}$ are Legendrian equivalent. Therefore we have that $\tilde{\pi}\circ i$ and $\tilde{\pi}\circ C'|_{\tilde{\mathbb{L}}}$ are Legendrian equivalent. ■

Theorem 11.7 *Let $\tilde{\pi}\circ i : (\tilde{\mathbb{L}},0) \to (J^1(\mathbb{R}^n,\mathbb{R}),0) \to (\mathbb{R}^n\times\mathbb{R},0)$ be a reticular Legendrian map. If $\tilde{\pi}\circ i$ is infinitesimally stable then $\tilde{\pi}\circ i$ is $(n+3)$-determined.*

Proof. It is enough to prove $\tilde{\pi}\circ i$ is $(n+2)$-determined of Definition 11.4 (3) and this is proved by an analogous method of Theorem 11.3. We write a sketch of the proof. Let $C \in C^{\theta}_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ be an extension of i. Then we may assume that P_C has the form

$$P_C = \{(Q, -\frac{\partial H}{\partial Q}(Q,p); H(Q,p) - \langle\frac{\partial H}{\partial Q}(Q,p), p\rangle, -\frac{\partial H}{\partial p}(Q,p), p)\}$$

for some function germ $H(Q,p) \in \mathfrak{M}(2n)^2$. Then $F(x,y,q,z) = H_0(x,y) + \langle y,q\rangle - z \in \mathfrak{M}(r; n+n+1)$ is a generating family of $\tilde{\pi}\circ i$, where $H_0(x,y) = H(x,0,y) \in \mathfrak{M}(r;n)^2$. Then F is a reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimally stable unfolding of $H_0(x,y)$. It follows that H_0 is reticular $\mathcal{K}$-$(n+2)$-determined. Let $C' \in C^{\theta}_Z(J^1(\mathbb{R}^n,\mathbb{R}),0)$ satisfying $j^{n+2}C(0) = j^{n+2}C'(0)$ be given. There exists a function germ $H'(Q,p) \in \mathfrak{M}(2n)$ such that

$$P_{C'} = \{(Q, -\frac{\partial H'}{\partial Q}(Q,p); H'(Q,p) - \langle\frac{\partial H'}{\partial Q}(Q,p), p\rangle, -\frac{\partial H'}{\partial p}(Q,p), p)\}.$$

Since $H = z - qp$ on P_C and $H' = z - qp$ on $P_{C'}$, we have that $j^{n+2}H_0(0) = j^{n+2}H_0'(0)$, where $H_0'(x,y) = H'(x,0,y) \in \mathfrak{M}(r;n)^2$. The function germ $G(x,y,q,z) = H_0'(x,y) + \langle y,q\rangle - z \in \mathfrak{M}(r; n+n+1)$ is a generating family of $\tilde{\pi}\circ C'|_{\tilde{\mathbb{L}}}$. Then there exist $\Phi(x,y) \in \mathcal{B}(r;n)$ and an unit $a \in \mathcal{E}(r;n)$ such that $H_0 = a\cdot H_0'\circ\Phi$. We set $G'(x,y,q,z) = a(x,y)G(\Phi(x,y),q,z) \in$

$\mathfrak{M}(r; n+n+1)$. Then G and G' are reticular $\mathcal{P}$-$\mathcal{K}$-equivalent and F and G' are reticular $\mathcal{P}$-$\mathcal{K}$-infinitesimal stable unfoldings of $H_0(x,y)$. It follows that F and G are reticular $\mathcal{P}$-$\mathcal{K}$-equivalent. Therefore $\tilde{\pi} \circ i$ and $\tilde{\pi} \circ C'|_{\tilde{\mathbb{L}}}$ are Legendrian equivalent. ■

In order to prove Theorem 12.5 and Theorem 13.4, we require the following lemmas:

Lemma 11.8 (cf., [2, p.53 Lemma 4.6]) *Let V, W be smooth manifolds with Q a submanifold of W, F be a topological space. We equip $C^\infty(V,W)$ with the Whitney C^∞-topology. Let $j : F \to C^\infty(V,W)$ be a map (not necessary continuous).* We suppose that: For each $f \in F$, there exist a manifold E, $e_0 \in E$, and a continuous map $\phi : E \to F$, $\phi(e_0) = f$ such that the map $\Phi : E \times V \to W$, $\Phi(e,x) = j(\phi(e))(x)$, is smooth and transversal to Q. Then the set

$$T = \{f \in F \mid j(f) \text{ is transversal to } Q\}$$

is dense in F.

Lemma 11.9 *Let V, W be smooth manifolds with Q a submanifold of W, and $K \subset Q$ be a compact set. Then the set*

$$T_K = \{f \in C^\infty(V,W) \mid j^l f \textit{ is transversal to } Q \textit{ on } K\}$$

is open in $C^\infty(V,W)$ with Whitney C^{l+1}-topology (hence Whitney C^∞-topology).

12 Genericity of reticular Lagrangian maps

Let $J^l(2n,2n)$ be the set of l-jets of map germs from $(T^*\mathbb{R}^n, 0)$ to $(T^*\mathbb{R}^n, 0)$ and $S^l(n)$ be the Lie group in $J^l(2n,2n)$ consists of l-jets of symplectic diffeomorphism germs on $(T^*\mathbb{R}^n, 0)$.

We consider the Lie group $L^l(2n) \times L^l(2n)$ acts on $J^l(2n,2n)$ as coordinate changes of the source and target spaces. We also consider the Lie subgroup $rLa^l(n)$ of $L^l(2n) \times L^l(2n)$ consists of l-jets of reticular diffeomorphisms on the source space and l-jets of Lagrangian equivalences of π:

$$rLa^l(n) = \{(j^l\phi(0), j^l\Theta(0)) \in L^l(2n) \times L^l(2n) \mid \phi \text{ is a reticular diffeomorphism on } (T^*\mathbb{R}^n, 0), \Theta \text{ is a Lagrangian equivalence of } \pi\}.$$

The group $rLa^l(n)$ acts on $J^l(2n,2n)$ and $S^l(n)$ is invariant under this action. Let S be a symplectic diffeomorphism germ on $(T^*\mathbb{R}^n, 0)$ and set $z = j^l S(0)$. We denote the orbit $rLa^l(n) \cdot z$ by $[z]$. Then

$$[z] = \{j^l S'(0) \in S^l(n) \mid \pi \circ i \text{ and } \pi \circ S'|_{\mathbb{L}} \text{ are Lagrangian equivalent}\}.$$

In this section we denote by X_f the Hamiltonian vector field on $(T^*\mathbb{R}^n, 0)$ for a function germ f on $(T^*\mathbb{R}^n, 0)$. That is

$$X_f = \sum_{j=1}^n \left(\frac{\partial f}{\partial p_j}\frac{\partial}{\partial q_j} - \frac{\partial f}{\partial q_j}\frac{\partial}{\partial p_j}\right).$$

We denote by VI_S the vector space consists of infinitesimal symplectic transformations of S and denote by VI_S^0 the subspace of VI_S consists of germs which vanishes on 0. We denote by $VL_{T^*\mathbb{R}^n}$ by the vector space consists of infinitesimal Lagrangian equivalences of π and denote by $VL^0_{T^*\mathbb{R}^n}$ by the subspace of $VL_{T^*\mathbb{R}^n}$ consists of germs which vanishes at 0.

We denote by $V^0_{\mathbb{L}}$ the vector space consists of infinitesimal reticular diffeomorphisms on $(T^*\mathbb{R}^n, 0)$ which vanishes at 0:

$$V^0_{\mathbb{L}} = \{\xi \in X(T^*\mathbb{R}^n, 0) \mid \xi \text{ is tangent to } L^0_\sigma \text{ for all } \sigma \subset I_r,\ \xi(0) = 0\}.$$

From now on, we denote by $\mathcal{E}_{T^*\mathbb{R}^n}$ the ring of smooth function germs on $(T^*\mathbb{R}^n, 0)$ and denote by $\mathfrak{M}_{T^*\mathbb{R}^n}$ its maximal ideal. We also denote other notations analogously.

Lemma 12.1 (1) *A vector field germ v on $(T^*\mathbb{R}^n, 0)$ along S is belongs to VI_S if and only if there exists a function germ f on $(T^*\mathbb{R}^n, 0)$ such that $v = X_f \circ S$,*
(2) *A vector field germ η on $(T^*\mathbb{R}^n, 0)$ is belongs to $VL_{T^*\mathbb{R}^n}$ if and only if there exists a fiber preserving function germ H on $(T^*\mathbb{R}^n, 0)$ such that $\eta = X_H$.*
(3) *A vector field germ ξ on $(T^*\mathbb{R}^n, 0)$ is belongs to $V^0_{\mathbb{L}}$ if and only if there exists a function germ $g \in B_0$ such that $\xi = X_g$, where $B_0 = \langle q_1p_1, \ldots, q_rp_r\rangle_{\mathcal{E}_{T^*\mathbb{R}^n}} + \mathfrak{M}_{T^*\mathbb{R}^n}\langle q_{r+1}, \ldots, q_n\rangle$ is a submodule of $\mathcal{E}_{T^*\mathbb{R}^n}$.*

By this lemma we have that:

$$VI^0_S = \{v : (T^*\mathbb{R}^n, 0) \to (T(T^*\mathbb{R}^n), 0) \mid v = X_f \circ S \text{ for some } f \in \mathfrak{M}^2_{T^*\mathbb{R}^n}\},$$

$$VL^0_{T^*\mathbb{R}^n} = \{\eta \in X(T^*\mathbb{R}^n, 0) \mid \eta = X_H \text{ for some fiver preserving function germ } H \in \mathfrak{M}^2_{T^*\mathbb{R}^n}\},$$

$$V^0_{\mathbb{L}} = \{\xi \in X(T^*\mathbb{R}^n, 0) \mid \xi = X_g \text{ for some } g \in B_0\}.$$

We define the homomorphism $tS : VI^0_{\mathbb{L}} \to VI^0_S$ by $tS(v) = S_*v$ and define the homomorphism $wS : VL^0_{T^*\mathbb{R}^n} \to VI^0_S$ by $wS(\eta) = \eta \circ S$.

Lemma 12.2 *Let S be a symplectic diffeomorphism germ on $(T^*\mathbb{R}^n, 0)$ and set $z = j^lS(0)$. Then*

$$T_z(rLa^l(n) \cdot z) = \pi_l(tS(V^0_{\mathbb{L}}) + wS(VL^0_{T^*\mathbb{R}^n})).$$

We denote VI^l_S the subspace of VI_S consists of infinitesimal symplectic transformation germs of S whose l-jets are 0:

$$VI^l_S = \{v \in VI_S \mid j^lv(0) = 0\}.$$

We consider the surjective projection $\pi_l : VI_S \to T_z(S^l(n))$. Since $(j^lS)_*(\frac{\partial}{\partial q_j}) = \pi_l(S_*\frac{\partial}{\partial q_j})$, $(j^lS)_*(\frac{\partial}{\partial p_j}) = \pi_l(S_*\frac{\partial}{\partial p_j})$, it follows that j^lS is transversal to $[z]$ if and only if $(j^lS)_*(T_0(T^*\mathbb{R}^n))$ $+T_z[z] = T_z(S^l(n))$ and this holds if and only if

$$(\pi_l)^{-1}((j^lS)_*(T_0(T^*\mathbb{R}^n)) + tS(V^0_{\mathbb{L}}) + wS(VL^0_{T^*\mathbb{R}^n})) + VI^{l+1}_S = VI_S$$

and this holds if and only if

$$tS(V^0_{\mathbb{L}}) + wS(VL_{T^*\mathbb{R}^n})) + VI^{l+1}_S = VI_S.$$

Let N, M be $2n$ dimensional symplectic manifolds. We denote $S(N,M)$ the space of symplectic embeddings from N to M with the topology induced from the Whitney C^∞-topology of $C^\infty(N,M)$. We define that

$$J^l_S(N,M) = \{j^l S(u_0) \in J^l(N,M) \mid \\ S : (N,u_0) \to M \text{ is a symplectic embedding germ, } u_0 \in N\}.$$

Proposition 12.3 *$S(N,M)$ is a Baire space.*

This is proved by an analogous method of the assertion that $C^\infty(N,M)$ is a Baire space (cf., [2, p.44 Proposition 3.3]).

Theorem 12.4 (Symplectic transversality theorem) *Let N, M be $2n$ dimensional symplectic manifolds. Let $Q_j, j = 1, 2, \ldots$ be submanifolds of $J^l_S(N,M)$. Then the set*

$$T = \{S \in S(N,M) \mid j^l S \text{ is transversal to } Q_j \text{ for all } j \in \mathbb{N}\}$$

is residual set in $S(N,M)$. In particular T is dense.

Proof. We apply for Lemma 11.8 that $V = N, W = J^l_S(N,M)$, and $F = S(N,M)$. We reduce our assertion to local situations by choosing a countable covering of Q_j by sufficiently small compact sets $K_{j,k}$'s. Then the sets $T_{j,k} = \{S \in S(N,M) \mid j^l S$ is transversal to Q_j at $K_{j,k}\}$ are open set by Lemma 11.8 and we have that $T = \cap T_{j,k}$ We fix a symplectic embedding $S \in S(N,M)$. For each $u_0 \in N$ there exist local symplectic coordinate systems of N around u_0 and M around $S(u_0)$ such that S is given by $(q,p) \mapsto (q,p)$ around 0.

For each j,k we take E by a sufficiently small neighborhood of 0 in $P(2n,1;l+1)$ and take a smooth function $\rho : T^*\mathbb{R}^n \to [0,1]$ such that ρ is 1 on a neighborhood of 0 and zero outside a compact set, where $P(2n,1;l+1)$ is the set of not higher than $(l+1)$-degree polynomials on $2n$ variables.

For each $H \in E$ we define $H'(Q,p) = \rho(Q,p)H(Q,p) - \langle Q,p\rangle$ and $\psi_{H'}(Q,p) = (Q, -\frac{\partial H'}{\partial Q}(Q,p))$ for $(Q,p) \in T^*\mathbb{R}^n$ around 0. Then there exists a neighbourhood U of 0 in $T^*\mathbb{R}^n$ such that $\psi_{H'}$ is a embedding on U and equal to the identity map outside a compact set for any $H \in E$. Therefore there exists a neighbourhood U' of 0 in $T^*\mathbb{R}^n$ such that the map $E \to C^\infty(U',U)$, $H \mapsto (\psi_H^{-1})|_{U'}$ is well defined and continuous. Each $(\psi_H^{-1})|_{U'}$ is equal the identity map outside a compact set around 0. We set that

$$\phi(H)(Q,P) = (-\frac{\partial H'}{\partial p}(Q,p), p) \circ (\psi_{H'})^{-1}(Q,P) \text{ for } (Q,P) \in U'.$$

Then $\phi(H)$ is a symplectic diffeomorphism around u_0 which has the canonical relation with the generating function $H'(Q,p)$ and equal to S outside a compact set. It follows that the source space of $\phi(H)$ may be extended naturally to $E \times N$. We also denote this by $\phi_H \in S(N,M)$. Then the map

$$\Phi : E \times N \to J^l_S(N,M), \Phi(H,q,p) = j^l(\phi_H)(q,p)$$

is a submersion around $(0,u_0)$ Therefore Φ is transversal to $K_{j,k}$. So we have the result. ■

For $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ we define the continuous map $j^l_0\tilde{S} : U \to S^l(n)$ by w to the l-jet of $\tilde{S}_w$ at 0. We remark that $j^l\tilde{S}_w(0)$ has the form $j^l\tilde{S}_w(0) = (w, \tilde{S}(w), j^l_0\tilde{S}_w(0))$.

Theorem 12.5 *Let U be a neighborhood of 0 in $T^*\mathbb{R}^n$, and let $Q_1, Q_2, \dots$ are submanifolds of $S^l(n)$. Then the set $L_U = \{(q,p) \in U | q = p_1 = \cdots = p_r = 0\}$ and*

$$T = \{\tilde{S} \in S(U, T^*\mathbb{R}^n) | j^l_0\tilde{S} \textit{ is transversal to } Q_j \textit{ on } L_U \textit{ for all } j\}$$

is a residual set in $S(U, T^*\mathbb{R}^n)$.

Proof. We set $Q'_j = L_U \times T^*\mathbb{R}^n \times Q_j \subset J^l_S(U, T^*\mathbb{R}^n)$. We choose a countable covering of Q'_j by sufficiently small compact sets $K_{j,k}{}'s$ for all j. We apply $N = V, M = T^*\mathbb{R}^n$ for Theorem 12.4. We have that

$$T = \{\tilde{S} \in S(U, T^*\mathbb{R}^n) | j^l\tilde{S} \text{ is transversal to } Q'_j \text{ on } K_{j,k} \text{ for all } j,k\}.$$

It follows that T is a residual set in $S(U, T^*\mathbb{R}^n)$. ■

Theorem 12.6 *Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a reticular Lagrangian map. Let S be an extension of i and $l \geq n+1$. Let $B = \langle q_1p_1, \dots, q_rp_r, q_{r+1}, \dots, q_n\rangle_{\mathcal{E}_{T^*\mathbb{R}^n}}$ be a submodule of $\mathcal{E}_{T^*\mathbb{R}^n}$. Then the followings are equivalent:*
(s) *$\pi \circ i$ is stable.*
(t) *$j^l_0 S$ is transversal to $[j^l_0 S(0)]$ at 0.*
(a') *$\mathcal{E}_{T^*\mathbb{R}^n}/(B + \mathfrak{M}^{l+2}_{T^*\mathbb{R}^n})$ is generated by $1, p_1 \circ S, \dots, p_n \circ S$ as $\mathcal{E}_q$-module via $\pi \circ S$.*
(a") *$\mathcal{E}_{T^*\mathbb{R}^n}/B$ is generated by $1, p_1 \circ S, \dots, p_n \circ S$ as $\mathcal{E}_q$-module via $\pi \circ S$.*
(is) *$\pi \circ i$ is infinitesimally stable.*

Proof. (s)$\Rightarrow$(t). Let $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ be a representative of S. By theorem 12.5 there exists a symplectic embedding $\tilde{S'}$ around $\tilde{S}$ such that $j^l_0\tilde{S'}$ is transversal to $[j^l_0 S(0)]$ at $w = (0, \dots, 0, p_{r+1}, \dots, p_n) \in U$. Since $\pi \circ i$ is stable, $\pi \circ i$ and $\pi \circ \tilde{S'_w}|_{\mathbb{L}}$ are Lagrangian equivalent. This means that $[j^l_0\tilde{S'_w}(0)] = [j^l_0 S(0)]$ and hence $j^l_0 S$ is transversal to $[j^l_0 S(0)]$ at 0.
(t)$\Leftrightarrow$(a'). By Lemma 12.1, we have that the condition (t) is equivalent to the condition: For any function germ f on $(T^*\mathbb{R}^n, 0)$, there exist a fiber preserving function germ H on $(T^*\mathbb{R}^n, 0)$ and a function germ $g \in B$ such that $\pi_l(X_f \circ S) = \pi_l(S_*X_g + X_H \circ S)$. This is equivalent to the condition: For any function germ f on $(T^*\mathbb{R}^n, 0)$, there exist a fiber preserving function germ H on $(T^*\mathbb{R}^n, 0)$ and a function germ $g \in B$ such that $f \circ S - g - H \circ S \in \mathfrak{M}^{l+2}_{T^*\mathbb{R}^n}$. This is equivalent to the condition: For any function germ f on $(T^*\mathbb{R}^n, 0)$, there exist a fiber preserving function germ H on $(T^*\mathbb{R}^n, 0)$ such that $f - H \circ S \in B + \mathfrak{M}^{l+2}_{T^*\mathbb{R}^n}$. This is equivalent to (a').
(a')$\Leftrightarrow$(a"). We need only to prove (a')$\Rightarrow$(a"). By Margrange preparation theorem, the condition (a') is equivalent that $\mathcal{E}_{T^*\mathbb{R}^n}/\ ((\pi \circ S)^*\mathfrak{M}_{\mathbb{R}^n}\mathcal{E}_{T^*\mathbb{R}^n} + B + \mathfrak{M}^{l+2}_{T^*\mathbb{R}^n})$ is generated by $1, p_1 \circ S, \dots, p_n \circ S$ over $\mathbb{R}$. This means that

$$\mathfrak{M}^{n+1}_{T^*\mathbb{R}^n} \subset (\pi \circ S)^*\mathfrak{M}_{\mathbb{R}^n}\mathcal{E}_{T^*\mathbb{R}^n} + B + \mathfrak{M}^{l+2}_{T^*\mathbb{R}^n}.$$

Since $\mathfrak{M}^{l+2}_{T^*\mathbb{R}^n} \subset \mathfrak{M}^{n+2}_{T^*\mathbb{R}^n}$, it follows that

$$\mathfrak{M}^{n+1}_{T^*\mathbb{R}^n} \subset (\pi \circ S)^*\mathfrak{M}_{\mathbb{R}^n}\mathcal{E}_{T^*\mathbb{R}^n} + B + \mathfrak{M}^{n+2}_{T^*\mathbb{R}^n}.$$

Therefore we have that

$$\mathfrak{M}^{n+1}_{T^*\mathbb{R}^n} \subset (\pi \circ S)^*\mathfrak{M}_{\mathbb{R}^n}\mathcal{E}_{T^*\mathbb{R}^n} + B.$$

This means (a").
(a")$\Leftrightarrow$(is). The condition (a") is equivalent to the condition (a) in Theorem 10.1 and this is

equivalent to (is).
(t)&(is)⇒(s). Since $j^l_0 S$ is transversal to $[j^l_0 S(0)]$, it follows that there exists a representative $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ of S and a neighbourhood $W_{\tilde{S}}$ of $\tilde{S}$ such that for any $\tilde{S}' \in W_{\tilde{S}}$ there exists $w \in U$ such that $j^l_0 \tilde{S}'$ is transversal to $[j^l_0 S(0)]$ at w. Since $j^l_0 \tilde{S'_w}(0) \in [j^l_0 S(0)]$, it follows that there exists a symplectic embedding germ S'' on $(T^*\mathbb{R}^n, 0)$ such that $\pi \circ i$ and $\pi \circ S''|_{\mathbb{L}}$ is Lagrangian equivalent and $j^l_0 S''(0) = j^l_0 \tilde{S'_w}(0)$. Since $\pi \circ i$ is infinitesimally stable, it follows that $\pi \circ i$ is l-determined by Theorem 11.3. Therefore we have that $\pi \circ S''|_{\mathbb{L}}$ is also l-determined. It follows that $\pi \circ S''|_{\mathbb{L}}$ and $\pi \circ \tilde{S'_w}|_{\mathbb{L}}$ is Lagrangian equivalent. This means that $\pi \circ i$ is stable. ■

Let S be a symplectic diffeomorphism germ on $(T^*\mathbb{R}^n, 0)$, $F(x, y, q) \in \mathfrak{M}(r; k+n)^2$, and $f(x, y) \in \mathfrak{M}(r; k)^2$. We denote $[S], [F], [f]$ by the equivalence classes of S, F, f under the Lagrangian equivalence, the stably reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalence, and the stably reticular $\mathcal{R}$-equivalence respectively. Then the following holds:

Lemma 12.7 *Let $\pi \circ i_j : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a stable reticular Lagrangian map with a generating family $F_j(x, y, q) \in \mathfrak{M}(r; k_j + n)^2$, and S_j be an extension of i_j for $j = 1, 2$. Then $[S_1] = [S_2]$ if and only if $[F_1] = [F_2]$ and this holds if and only if $[F_1|_{q=0}] = [F_2|_{q=0}]$.*

Proof. $[S_1] = [S_2]$ if and only if reticular Lagrangian maps $\pi \circ i_1$ and $\pi \circ i_2$ are Lagrangian equivalent by Lemma 11.2, and this holds if and only if $[F_1] = [F_2]$ by Theorem 6.2(3), and if this holds then $[F_1|_{q=0}] = [F_2|_{q=0}]$. Conversely suppose that $[F_1|_{q=0}] = [F_2|_{q=0}]$. Since F_1 and F_2 are reticular $\mathcal{P}$-$\mathcal{R}^+$-stable unfoldings of $[F_1|_{q=0}]$ and $[F_2|_{q=0}]$ respectively and $F_1|_{q=0}$ and $F_2|_{q=0}$ are stably reticular $\mathcal{R}$-equivalent. it follows that $[F_1] = [F_2]$. ■

Corollary 12.8 *Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a stable reticular Lagrangian map. Then there exist a neighborhood U of 0 in $T^*\mathbb{R}^n$ and $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ with $i = \tilde{S}_0|_{\mathbb{L}}$ (that is, $\tilde{S}$ is a representative of a extension of i) such that reticular Lagrangian maps $\pi \circ \tilde{S}_w|_{\mathbb{L}}$ are stable for all $w \in U$*

Proof. By Theorem 12.6 (a'), the stability of reticular Lagrangian maps are determined by the $(n+1)$-jets of $\pi \circ \tilde{S}_w$ for $w \in U$. Therefore we have the result by shrinking U if necessary. ■

Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a stable reticular Lagrangian map. We say that $\pi \circ i$ is *simple* if there exist a neighborhood U of 0 in $T^*\mathbb{R}^n$ and $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ such that $i = \tilde{S}_0|_{\mathbb{L}}$ and $\{\tilde{S}_w | w \in U\}$ is covered by finite orbits $[S_1], \ldots, [S_m]$ for symplectic diffeomorphism germs $S_1, \ldots, S_m$ on $(T^*\mathbb{R}^n, 0)$.

Lemma 12.9 *Let $\pi \circ i : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ be a stable reticular Lagrangian map. Then $\pi \circ i$ is simple if and only if there exist a neighborhood U_z of $z = j^{n+1}_0 S(0)$ in $S^{n+1}(n)$ and $z_1, \ldots, z_m \in S^{n+1}(n)$ such that $U_z \subset [z_1] \cup \cdots \cup [z_m]$.*

Proof. Suppose that $\pi \circ i$ is simple. Then there exists a representative $\tilde{S} : U \to T^*\mathbb{R}^n$ of an extension of i and symplectic diffeomorphism germs $[S_1], \ldots, [S_m]$ on $(T^*\mathbb{R}^n, 0)$ such that

$$\{\tilde{S}_w | w \in U\} \subset [S_1] \cup \cdots \cup [S_m]. \tag{1}$$

Since $\pi \circ i$ is stable, it follows that $j^{n+1}_0 \tilde{S}$ is transversal to $[z]$ at 0 by Theorem 7.1. This means that there exists a neighbourhood U_z of z in $S^l(n)$ such that $U_z \subset \cup_{w \in U} [j^{n+1}_0 \tilde{S}(w)]$. It follows that $U_z \subset [j^{n+1} S_1(0)] \cup \cdots \cup [j^{n+1} S_m(0)]$.

Conversely suppose that there exist a neighbourhood U_z of z in $S^{n+1}(n)$ and $z_1, \dots, z_m \in S^{n+1}(n)$ such that $U_z \subset [z_1] \cup \cdots \cup [z_m]$. Since the map $j_0^{n+1}\tilde{S} : U \to S^{n+1}(n)$ is continuous, there exists a neighbourhood U' of 0 in U such that $j_0^{n+1}\tilde{S}(w) \in U_z$ for any $w \in U'$. Then we have that $\cup_{w \in U'} j_0^{n+1}\tilde{S}(w) \subset [z_1] \cup \cdots \cup [z_m]$. Choose symplectic diffeomorphism germs $S_1, \dots, S_m$ on $(T^*\mathbb{R}^n, 0)$ such that $j^{n+1}S_j(0) = z_j$ for $j = 1, \dots, m$. By Corollary 12.8, we may assume that each $\pi \circ S_j|_{\mathbb{L}}$ is stable, therefore $(n+1)$-determined. For any $w \in U'$ we have that there exists $j \in \{1, \dots, m\}$ such that $j_0^{n+1}\tilde{S}_w(0) \in [j^{n+1}S_j(0)]$. It follows that reticular Lagrangian maps $\pi \circ \tilde{S}_w|_{\mathbb{L}}$ and $\pi \circ S_j|_{\mathbb{L}}$ are Lagrangian equivalent. Therefore $\tilde{S}_w \in [S_j]$. We have (1). ■

Lemma 12.10 *A stable reticular Lagrangian map $\pi \circ i$ is simple if and only if for a generating family $F(x, y, q) \in \mathfrak{M}(r; k+n)^2$ of $\pi \circ i$, the function germ $F(x, y, 0) \in \mathfrak{M}(r; k)^2$ is $\mathcal{R}$-simple singularity.*

Proof. Let S be an extension of i and $\tilde{S} \in S(U, T^*\mathbb{R}^n)$ be a representative of S

Suppose that $F(x, y, 0)$ is $\mathcal{R}$-simple. The simplicity of reticular Lagrangian maps is invariant under Lagrangian equivalences, we may assume that the map germ $(Q, P) \mapsto (Q, p \circ S(Q, P))$ is a diffeomorphism germ on $(\mathbb{R}^{2n}, 0)$. We consider a symplectic diffeomorphism germ $\tilde{S}_w$ on $(\mathbb{R}^{2n}, 0)$ for $w \in U$ near 0. Then there exists a function germ $H_w(Q, p) \in \mathfrak{M}(2n)^2$ such that the canonical relation P_w associated with $\tilde{S}_w$ has the form:

$$P_w = \{(Q, -\frac{\partial H_w}{\partial Q}(Q, p), -\frac{\partial H_w}{\partial p}(Q, p), p) \in (T^*\mathbb{R}^n \times T^*\mathbb{R}^n, (0, 0))\}.$$

Then the function germ $F_w(x, y, q) = H'_w(x, y) + \langle y, q\rangle \in \mathfrak{M}(r; n+n)^2$ is a generating family of $\pi \circ \tilde{S}_w|_{\mathbb{L}}$, where $H'_w \in \mathfrak{M}(r; n)^2$ is defined by $H'_w(x, y) = H_w(x, 0, y)$. Since F_0 is a generating family of $\pi \circ i$, we have that $F_0(x, y, 0) (= H'_0(x, y))$ is stably $\mathcal{R}$-equivalent to $F(x, y, 0)$. Therefore we have that H'_0 is $\mathcal{R}$-simple. Then there exists $f_1, \dots, f_m \in \mathfrak{M}(r; n)$ and a neighbourhood V of $j^{n+2}H'_0(0)$ in $J^{n+2}(r+n, 1)$ such that $V \subset [j^{n+2}f_1(0)] \cup \cdots \cup [j^{n+2}f_m(0)]$.

Since the $(n+2)$-jet of H'_S is determined by the $(n+1)$-jet of S, there exists a neighbourhood U' of 0 in U such that the map germ

$$U' \to J^{n+2}(r+n, 1), \ w \mapsto j^{n+2}H'_w(0)$$

is well defined and continuous.

Let U'' be the inverse image of V by the above map. Then for any $w \in U''$ the reticular Lagrangian map $\pi \circ \tilde{S}_w|_{\mathbb{L}}$ has a generating family which is reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent to $f_j(x, y) + \langle y, q\rangle \in \mathfrak{M}(r; n+n)^2$ for some j because $\pi \circ \tilde{S}_w|_{\mathbb{L}}$ is stable by Corollary 12.8 and hence $(n+1)$-determined. It follow that $\{\tilde{S}_w | w \in U''\} \subset [S_1] \cup \cdots \cup [S_m]$, where S_j is an extension of a reticular Lagrangian embedding which defines a reticular Lagrangian map with the generating family $f_j(x, y) + \langle y, q\rangle \in \mathfrak{M}(r; n+n)^2$. This means that $\pi \circ i$ is simple.

Conversely suppose that $\pi \circ i$ is simple. Let S_0 be an extension of i. we may assume that there exists a function germ $H_0(Q, p) \in \mathfrak{M}(2n)^2$ such that the canonical relation P_{S_0} associated with S_0 has the form:

$$P_{S_0} = \{(Q, -\frac{\partial H_0}{\partial Q}(Q, p), -\frac{\partial H_0}{\partial p}(Q, p), p) \in (T^*\mathbb{R}^n \times T^*\mathbb{R}^n, (0, 0))\}.$$

This means that for a function germ H around H_0 there exists a symplectic diffeomorphism germ S_H on $(T^*\mathbb{R}^n, 0)$ with the canonical relation P_{S_H} which has the form same as P_{S_0}.

Then the $(n+1)$-jet of S_H is determined by the $(n+2)$-jet of H. The function germ $H_0'(x,0,y)+\langle y,q\rangle \in \mathfrak{M}(r;n+n)^2$ is a generating family of $\pi\circ i$, where $H_0' \in \mathfrak{M}(r;n)^2$ is defined by $H_0'(x,y) = H_0(x,0,y)$. Then there exists a quadratic form $T(Q_{r+1},\dots,Q_n,p_1,\dots,p_n)$ such that the function germ $H_0''(Q,p) = H_0'(Q_1,\dots,Q_r,p)+T(Q_{r+1},\dots,Q_n,p)$ also defined the symplectic diffeomorphism $S_{H_0''}$. Then we have that $\pi\circ S_{H_0''}|_{\mathbb{L}} = \pi\circ i$. Therefore we may assume that $H_0'' = H_0$

We have that H_0' is stably reticular $\mathcal{R}$-equivalent to $F(x,y,0)$. Therefore we need only to prove that H_0' is a simple singularity.

There exists a neighbourhood V of $j^{n+2}H_0'(0)$ such that the map

$$V \to S^{n+1}(n),\ j^{n+2}H'(0) \mapsto j_0^{n+1}S_H(0),$$

where $H(Q,p) = H'(Q_1,\dots,Q_r,p)+T(Q_{r+1},\dots,Q_n,p)$ is well defined and continuous.

Since $\pi\circ i$ is simple, there exist finite symplectic diffeomorphism germs $S_1,\dots,S_m$ on $(T^*\mathbb{R}^n,0)$ and a neighbourhood U_z of $z = j^{n+1}S_0(0)$ in $S^{n+1}(n)$ such that $U_z \subset [S_1]\cup\cdots\cup[S_m]$ Let V' be the inverse image of U_z by the above map. Then we have that $V' \subset [z_1]\cup\cdots\cup[z_m]$, where $z_j = j^{n+2}H_j(0)$ for a function germ $H_j' \in \mathcal{E}(r;n)$ such that $j^{n+1}S_j(0) = j^{n+1}S_{H_j}(0)$. This means that H_0' is simple.

Theorem 12.11 *Let $r=0, n\leq 5$ or $r=1, n\leq 3$. Let U be a neighborhood of 0 in $T^*\mathbb{R}^n$. Then there exists a residual set $O \subset S(U,T^*\mathbb{R}^n)$ such that for any $\tilde{S}\in O$ and $w=(0,\dots,p_{r+1}^0,\dots,p_n^0)\in U$, the reticular Lagrangian map $\pi\circ\tilde{S}_w|_{\mathbb{L}}$ is stable.*

In the case $r=0,\ n\leq 5$. A reticular Legendrian map $\pi\circ\tilde{S}_w|_{\mathbb{L}}$ for any $\tilde{S}\in O$ and $w\in U$ has a generating family F which is a reticular $\mathcal{P}$-$\mathcal{R}^+$-stable unfolding of one of $A_2, A_3^\pm, A_4, A_5^\pm, A_6, D_4^\pm, D_4^\pm, D_6^\pm, E_6^\pm$, that is F is stably reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent to one of the following list:
$A_2 : F(y_1,q_1) = y_1^3+q_1y_1$,
$A_3^\pm : F(y_1,q_1,q_2) = \pm y_1^4+q_1y_1^2+q_2y_1$,
$A_4 : F(y_1,q_1,q_2,q_3) = y_1^5+q_1y_1^3+q_2y_1^2+q_3y_1$,
$A_5^\pm : F(y_1,q_1,q_2,q_3,q_4) = \pm y_1^6+q_1y_1^4+q_2y_1^3+q_3y_1^2+q_4y_1$,
$A_6 : F(y_1,q_1,q_2,q_3,q_4,q_5) = y_1^7+q_1y_1^5+q_2y_1^4+q_3y_1^3+q_4y_1^2+q_5y_1$,
$D_4^\pm : F(y_1,y_2,q_1,q_2,q_3) = y_1^2y_2\pm y_2^3+q_1y_2^2+q_2y_2+q_3y_1$,
$D_5^\pm : F(y_1,y_2,q_1,q_2,q_3,q_4) = y_1^2y_2\pm y_2^4+q_1y_2^3+q_2y_2^2+q_3y_2+q_4y_1$,
$D_6^\pm : F(y_1,y_2,q_1,q_2,q_3,q_4,q_5) = y_1^2y_2\pm y_2^5+q_1y_2^4+q_2y_2^3+q_3y_2^2+q_4y_2+q_5y_1$,
$E_6^\pm : F(y_1,y_2,q_1,q_2,q_3,q_4,q_5) = y_1^3\pm y_2^4+q_1y_1y_2^2+q_2y_1y_2+q_3y_2^2+q_4y_1+q_5y_2$.

In the case $r=1,\ n\leq 3$. A reticular Legendrian map $\pi\circ\tilde{S}_w|_{\mathbb{L}}$ for any $\tilde{S}\in O$ and $w\in U$ has a generating family which is a $\mathcal{P}$-$\mathcal{R}^+$-stable unfolding of one of $B_2^\pm, B_3^\pm, B_4^\pm, C_3^\pm, C_4^\pm, F_4^\pm$, that is F is stably reticular $\mathcal{P}$-$\mathcal{R}^+$-equivalent to one of the following list:
$B_2^\pm : F(x,q_1) = \pm x^2+q_1x$,
$B_3^\pm : F(x,q_1,q_2) = \pm x^3+q_1x^2+q_2x$,
$B_4^\pm : F(x,q_1,q_2,q_3) = \pm x^4+q_1x^3+q_2x^2+q_1x$,
$C_3^\pm : F(x,y,q_1,q_2) = \pm xy+y^3+q_1y^2+q_2y$,
$C_4^\pm : F(x,y,q_1,q_2,q_3) = \pm xy+y^4+q_1y^3+q_2y^2+q_3y$,
$F_4^\pm : F(x,y,q_1,q_2,q_3) = \pm x^2+y^3+q_1xy+q_2x+q_3y$.

Proof. We need only to prove the case $r=1,\ n\leq 3$. Let $F_X(x,y,q)\in\mathfrak{M}(r;k+n)^2$ be a reticular $\mathcal{P}$-$\mathcal{R}^+$-stable unfolding of singularity $X \in \mathfrak{M}(r;k)^2$ for

$$X = B_2^\pm, B_3^\pm, B_4^\pm, C_3^\pm, C_4^\pm, F_4^\pm.$$

Then other unfoldings are not stable since other singularities have reticular $\mathcal{R}^+$-codimension > 3. We choose stable reticular Lagrangian maps $\pi \circ i_X : (\mathbb{L}, 0) \to (T^*\mathbb{R}^n, 0) \to (\mathbb{R}^n, 0)$ with generating family F_X and S_X be and extension of i_X for above list. We set $L_U = \{(q,p) \in U | q = p_1 = 0\}$ and define that

$$O' = \{\tilde{S} \in S(U, T^*\mathbb{R}^n) \mid j_0^{n+1}\tilde{S} \text{ is transversal to } [j^{n+1}S_X(0)] \text{ on } L_U \text{ for all } X\}.$$

Then O' is a residual set. We set

$$Y = \{j^{n+1}S(0) \in S^{n+1}(n) \mid \text{the codimension of } [j^{n+1}S(0)] > 2n\}.$$

Then Y is an algebraic set in $S^{n+1}(n)$ by Theorem 12.6 (a'). Therefore we can define that

$$O'' = \{\tilde{S} \in S(U, T^*\mathbb{R}^n) \mid j_0^{n+1}\tilde{S} \text{ is transversal to } Y\}.$$

Then Y has codimension $> 2n$ because all element in Y is adjacent to one of the above list which are simple. Then we have that

$$O'' = \{\tilde{S} \in S(U, T^*\mathbb{R}^n) \mid j_0^{n+1}\tilde{S}(U) \cap Y = \emptyset\}.$$

We define $O = O' \cap O''$. Then O has the required condition.

In the case $r = 0, n \leq 5$. Set $X = A_2, A_3^\pm, A_4, A_5^\pm, A_6, D_4^\pm, D_4^\pm, D_6^\pm, E_6^\pm$ and $Y = \{j^{n+1}S(0) \in S^{n+1}(n) \mid \text{the codimension of } [j^{n+1}S(0)] > 2n\}$. Then we have that the codimension of Y in $S^{n+1}(n)$ is higher than $2n$ and the assertion is proved by the parallel method of the above case. ■



Figure 2: the caustic $B_3^{\pm}$



Figure 3: the caustics $C_3^{\pm}$



Figure 4: the caustics $B_4^\pm$



Figure 5: the caustics $C_4^\pm$



Figure 6: the caustics $F_4^{\pm}$

13 Genericity of reticular Legendrian maps

Let $J^l(2n+1, 2n+1)$ be the set of l-jets of map germs from $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ to $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ and $C^l(n)$ be the Lie group in $J^l(2n+1, 2n+1)$ consists of l-jets of contact diffeomorphism germs on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$. We consider the Lie subgroup $rLe^l(n)$ of $L^l(2n+1) \times L^l(2n+1)$ consists of l-jets of reticular diffeomorphisms on the source space and l-jets of Legendrian equivalences of $\tilde{\pi}$:

$$rLe^l(n) = \{(j^l\phi(0), j^l\Theta(0)) \in L^l(2n+1) \times L^l(2n+1) \mid \phi \text{ is a reticular}$$
$$\text{diffeomorphism on } (J^1(\mathbb{R}^n, \mathbb{R}), 0), \Theta \text{ is a Legendrian equivalence of } \tilde{\pi}\}.$$

The group $rLe^l(n)$ acts on $J^l(2n+1, 2n+1)$ and $C^l(n)$ is invariant under this action. Let C be a contact diffeomorphism germ on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ and set $z = j^lC(0)$. We denote the orbit $rLe^l(n) \cdot z$ by $[z]$. Then

$$[z] = \{j^lC'(0) \in C^l(n) \mid \tilde{\pi} \circ i \text{ and } \tilde{\pi} \circ C'|_{\tilde{\mathbb{L}}} \text{ are Legendrian equivalent}\}.$$

In this section we denote by X_f the Contact Hamiltonian vector field on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ for a function germ f on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$. That is

$$X_f = \sum_{j=1}^{n}(\frac{\partial f}{\partial q_j} + p_j \frac{\partial f}{\partial z})\frac{\partial}{\partial p_j} - \sum_{j=1}^{n} \frac{\partial f}{\partial p_j}\frac{\partial}{\partial q_j} + (f - \sum_{j=1}^{n} p_j \frac{\partial f}{\partial p_j})\frac{\partial}{\partial z}.$$

We denote by VI_C the vector space consists of infinitesimal contact transformation germs of C and denote by VI_C^0 the subspace of VI_C consists of germs which vanish on 0. We denote by $VL_{J^1(\mathbb{R}^n,\mathbb{R})}$ by the vector space consists of infinitesimal Legendrian equivalences on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ and denote by $VL^0_{J^1(\mathbb{R}^n,\mathbb{R})}$ by the subspace of $VL_{J^1(\mathbb{R}^n,\mathbb{R})}$ consists of germs which vanish at 0. We denote by $V^0_{\tilde{\mathbb{L}}}$ the vector space consists of infinitesimal reticular diffeomorphisms on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$ which vanishes at 0:

$$V^0_{\tilde{\mathbb{L}}} = \{\xi \in X(J^1(\mathbb{R}^n, \mathbb{R}), 0) \mid \xi \text{ is tangent to } \tilde{L}^0_\sigma \text{ for all } \sigma \subset I_r,\ \xi(0) = 0\}.$$

Lemma 13.1 *(1) A vector field germ v on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ along C is belongs to VI_C if and only if there exists a function germ f on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ such that $v = X_f \circ C$,*
(2) A vector field germ η on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ is belongs to $VL_{J^1(\mathbb{R}^n,\mathbb{R})}$ if and only if there exists a fiber preserving function germ H on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ such that $\eta = X_H$.
(3) A vector field germ ξ on $(J^1(\mathbb{R}^n,\mathbb{R}),0)$ is belongs to $V^0_{\tilde{\mathbb{L}}}$ if and only if there exists a function germ $g \in B'_0$ such that $\xi = X_g$, where $B'_0 = \langle q_1p_1,\dots,q_rp_r\rangle_{\mathcal{E}_{J^1(\mathbb{R}^n,\mathbb{R})}} + \mathfrak{M}_{J^1(\mathbb{R}^n,\mathbb{R})}\langle q_{r+1},\dots, q_n, z\rangle$ is a submodule of $\mathcal{E}_{J^1(\mathbb{R}^n,\mathbb{R})}$

By this lemma we have that:

$$VI^0_C = \{v : (J^1(\mathbb{R}^n,\mathbb{R}),0) \to (T(J^1(\mathbb{R}^n,\mathbb{R})),0) \mid v = X_f \circ C \text{ for some } f \in \mathfrak{M}^2_{J^1(\mathbb{R}^n,\mathbb{R})}\},$$

$$VL^0_{J^1(\mathbb{R}^n,\mathbb{R})} = \{\eta \in X(J^1(\mathbb{R}^n,\mathbb{R}),0) \mid \eta = X_H \\ \text{for some fiver preserving function germ } H \in \mathfrak{M}^2_{J^1(\mathbb{R}^n,\mathbb{R})}\},$$

$$V^0_{\tilde{\mathbb{L}}} = \{\xi \in X(J^1(\mathbb{R}^n,\mathbb{R}),0) \mid \xi = X_g \text{ for some } g \in B'_0\}.$$

We define the homomorphism $tC : VI^0_{\tilde{\mathbb{L}}} \to VI^0_C$ by $tC(v) = C_*v$ and define the homomorphism $wC : VL^0_{J^1(\mathbb{R}^n,\mathbb{R})} \to VI^0_C$ by $wC(\eta) = \eta \circ C$.

We denote VI^l_C the subspace of VI_C consists of infinitesimal contact transformation germs of C whose l-jets are 0:

$$VI^l_C = \{v \in VI_C \mid j^l v(0) = 0\}.$$

We have that $j^l_0 C$ is transversal to $[z]$ if and only if

$$tC(V^0_{\tilde{\mathbb{L}}}) + wC(VL_{J^1(\mathbb{R}^n,\mathbb{R})})) + VI^{l+1}_C = VI_C.$$

Let N, M be $(2n+1)$-dimensional contact manifolds. We denote $C(N,M)$ the space of contact embeddings from N to M with the topology induced from the Whitney C^∞-topology of $C^\infty(N,M)$ and define that

$$J^l_C(N,M) = \{j^l C(u_0) \in J^l(N,M) | C : (N,u_0) \to M \text{ is a contact embedding germ, } u_0 \in N\}.$$

Proposition 13.2 *$C(N,M)$ is a Baire space.*

Theorem 13.3 (Contact transversality theorem) *Let N, M be $(2n+1)$-dimensional contact manifolds. Let $Q_j, j = 1,2,\dots$ be submanifolds of $J^l_C(N,M)$. Then the set*

$$T = \{C \in C(N,M) \mid j^l C \text{ is transversal to } Q_j \text{ for all } j \in \mathbb{N}\}$$

is residual set in $C(N,M)$. In particular T is dense.

Theorem 13.4 *Let U be a neighborhood 0 in $J^1(\mathbb{R}^n, \mathbb{R})$, $Q_1, Q_2, \dots$ are submanifolds of $C^l(n)$. We define $L_U = \{(q, z, p) \in U | q = z = p_1 = \cdots = p_r = 0\}$ Then the set*

$$T = \{\tilde{C} \in C(U, J^1(\mathbb{R}^n, \mathbb{R})) | j_0^l \tilde{C} \text{ is transversal to } Q_j \text{ on } L_U \text{ for all } j\}$$

is a residual set in $C(U, J^1(\mathbb{R}^n, \mathbb{R}))$.

We have the following theorem which is proved by a parallel method of Theorem 12.6

Theorem 13.5 *Let $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ be a reticular Legendrian map. Let C be an extension of i and $l \geq n+2$. Let $B' = \langle q_1 p_1, \dots, q_r p_r, q_{r+1}, \dots, q_n, z \rangle_{\mathcal{E}_{J^1(\mathbb{R}^n, \mathbb{R})}}$ be a submodule of $\mathcal{E}_{J^1(\mathbb{R}^n, \mathbb{R})}$. Then the followings are equivalent:*
(s) *$\tilde{\pi} \circ i$ is stable.*
(t) *$j_0^l C$ is transversal to $[j_0^l C(0)]$.*
(a') *$\mathcal{E}_{J^1(\mathbb{R}^n, \mathbb{R})} / (B' + \mathfrak{M}_{J^1(\mathbb{R}^n, \mathbb{R})}^{l+2})$ is generated by $1, p_1 \circ C, \dots, p_n \circ C$ as $\mathcal{E}_{q,z}$-module via $\tilde{\pi} \circ C$.*
(a") *$\mathcal{E}_{J^1(\mathbb{R}^n, \mathbb{R})} / B'$ is generated by $1, p_1 \circ C, \dots, p_n \circ C$ as $\mathcal{E}_{q,z}$-module via $\tilde{\pi} \circ C$.*
(is) *$\tilde{\pi} \circ i$ is infinitesimally stable.*

For $\tilde{C} \in C(U, J^1(\mathbb{R}^n, \mathbb{R}))$ we define the continuous map $j_0^l \tilde{C} : U \to C^l(n)$ by w to the l-jet of $\tilde{C}_w$.

Let $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ be a stable reticular Legendrian map. We say that $\tilde{\pi} \circ i$ is *simple* if there exists a representative $\tilde{C} \in C(U, J^1(\mathbb{R}^n, \mathbb{R}))$ of a extension of i such that $\{\tilde{C}_w | w \in U\}$ is covered by finite orbits $[C_1], \dots, [C_m]$ for some contact diffeomorphism germs $C_1, \dots, C_m$ on $(J^1(\mathbb{R}^n, \mathbb{R}), 0)$.

By an analogous way of Lemma 12.9, Lemma 12.10 and Theorem 12.11 we have the followings:

Lemma 13.6 *Let $\tilde{\pi} \circ i : (\tilde{\mathbb{L}}, 0) \to (J^1(\mathbb{R}^n, \mathbb{R}), 0) \to (\mathbb{R}^n \times \mathbb{R}, 0)$ be a stable reticular Legendrian map. Then $\tilde{\pi} \circ i$ is simple if and only if there exist a neighbourhood U_z of $z = j_0^{n+3} C(0)$ in $C^{n+3}(n)$ and $z_1, \dots, z_m \in C^{n+3}(n)$ such that $U_z \subset [z_1] \cup \cdots \cup [z_m]$.*

Lemma 13.7 *A stable reticular Legendrian map $\tilde{\pi} \circ i$ is simple if and only if for a generating family $F(x, y, q, z) \in \mathfrak{M}(r; k+n+1)$ of $\tilde{\pi} \circ i$, $F(x, y, 0, 0) \in \mathfrak{M}(r; k)^2$ is $\mathcal{K}$-simple singularity.*

Theorem 13.8 *Let $r = 0, n \leq 6$ or $r = 1, n \leq 4$. Let U be a neighbourhood of 0 in $J^1(\mathbb{R}^n, \mathbb{R})$. Then there exists a residual set $O \subset C(U, J^1(\mathbb{R}^n, \mathbb{R}))$ such that for any $\tilde{C} \in O$ and $w = (0, \dots, 0, p_{r+1}, \dots, p_n) \in U$, the reticular Legendrian map $\tilde{\pi} \circ \tilde{C}_w|_{\tilde{\mathbb{L}}}$ is stable.*

In the case $r = 0, n \leq 6$. A reticular Legendrian map $\tilde{\pi} \circ \tilde{C}_w|_{\tilde{\mathbb{L}}}$ for any $\tilde{C} \in O$ and $w \in U$ has a generating family which is a reticular $\mathcal{P}$-$\mathcal{K}$-stable unfolding of one of $A_1, A_2, A_3, A_4, A_4, A_6, D_4^{\pm}, D_5, D_6^{\pm}, E_6$, that is F is stably reticular $\mathcal{P}$-$\mathcal{K}$-equivalent to one of the following list:
$A_2 : F(y_1, z) = y_1^2 + z,$
$A_2 : F(y_1, q_1, z) = y_1^3 + q_1 y_1 + z,$
$A_3 : F(y_1, q_1, q_2, z) = y_1^4 + q_1 y_1^2 + q_2 y_1 + z,$
$A_4 : F(y_1, q_1, q_2, q_3) = y_1^5 + q_1 y_1^3 + q_2 y_1^2 + q_3 y_1 + z,$
$A_5 : F(y_1, q_1, q_2, q_3, q_4) = y_1^6 + q_1 y_1^4 + q_2 y_1^3 + q_3 y_1^2 + q_4 y_1 + z,$
$A_6 : F(y_1, q_1, q_2, q_3, q_4, q_5, z) = y_1^7 + q_1 y_1^5 + q_2 y_1^4 + q_3 y_1^3 + q_4 y_1^2 + q_5 y_1 + z,$

$D_4^{\pm} : F(y_1, y_2, q_1, q_2, q_3, z) = y_1^2 y_2 \pm y_2^3 + q_1 y_2^2 + q_2 y_2 + q_3 y_1 + z,$
$D_5 : F(y_1, y_2, q_1, q_2, q_3, q_4, z) = y_1^2 y_2 + y_2^4 + q_1 y_2^3 + q_2 y_2^2 + q_3 y_2 + q_4 y_1 + z,$
$D_6^{\pm} : F(y_1, y_2, q_1, q_2, q_3, q_4, q_5, z) = y_1^2 y_2 \pm y_2^5 + q_1 y_2^4 + q_2 y_2^3 + q_3 y_2^2 + q_4 y_2 + q_5 y_1 + z,$
$E_6 : F(y_1, y_2, q_1, q_2, q_3, q_4, q_5, z) = y_1^3 + y_2^4 + q_1 y_1 y_2^2 + q_2 y_1 y_2 + q_3 y_2^2 + q_4 y_1 + q_5 y_2 + z.$

In the case $r = 1,\ n \leq 4$. A reticular Legendrian map $\tilde{\pi} \circ \tilde{C}_w|_{\tilde{\mathbb{L}}}$ for any $\tilde{C} \in O$ and $w \in U$ has a generating family which is a reticular $\mathcal{P}$-$\mathcal{K}$-stable unfolding of one of $B_2, B_3, B_4, C_3^{\pm}, C_4, F_4$, that is F is stably reticular $\mathcal{P}$-$\mathcal{K}$-equivalent to one of the following list:
$B_2 : F(x, q_1, z) = x^2 + q_1 x + z,$
$B_3 : F(x, q_1, q_2, z) = x^3 + q_1 x^2 + q_2 x + z,$
$B_4 : F(x, q_1, q_2, q_3, z) = x^4 + q_1 x^3 + q_2 x^2 + q_1 x + z,$
$C_3^{\pm} : F(x, y, q_1, q_2, z) = \pm xy + y^3 + q_1 y^2 + q_2 y + z,$
$C_4 : F(x, y, q_1, q_2, q_3, z) = xy + y^4 + q_1 y^3 + q_2 y^2 + q_3 y + z,$
$F_4 : F(x, y, q_1, q_2, q_3, z) = x^2 + y^3 + q_1 xy + q_2 x + q_3 y + z.$

References

[1] V.I.Arnold, S.M. Gusein-Zade & A.N. Varchenko, *Singularities of differential maps,* Vol.I, Birkhäuser, Basel, 1986.

[2] M.Golubitsky, V.Guillemin, *Stable Mappings and Their Singularities*, Graduate Text in Math. 14, Springer-Verlag, New York 1973.

[3] S.Izumiya G.Ishikawa, *Singularity Theory and their applications*, Kyouritsu shuppan, 1998 (in Japanese)

[4] S.Izumiya, *Semi-local Classification of Geometric Singularities for Hamilton-Jacobi Equations*, J. Differentials Geom., **118** (1995), pp.166-193

[5] Th:bröcker, *Differentiable Germs and Catastrophes*, London Math. Soc. Lecture Note ser., 17 Cambridge Univ. Press, 1975.

[6] S.Janeszko, *Generalized Luneburg canonical varieties and vector fields on quasicaustics*, J. Math. Phys. 31:4(1990), pp.997-1009.

[7] K.Jänich, *Caustics and catastrophes*, Math. Ann., 209(1974), pp.161-180.

[8] T.Tsukada, *Reticular Lagrangian, legendrian Singularities and their applications*, PhD Thesis, Hokkaido University, 1999.

[9] T.Tsukada, *Reticular Lagrangian Singularities*, The Asian J. of Math., vol 1, **3** (1997), pp.572-622.

[10] T.Tsukada, *Reticular Legendrian Singularities*, The Asian J. of Math., vol 5, **1** (2001), pp.109-127.

[11] G.Wassermann, *Stability of unfolding*, Lecture note in mathematics, 393.



Figure 7: the wavefronts B_2 and B_3



Figure 8: the wavefronts $C_3^{\pm}$

[12] G.Wassermann, *Stability of unfolding in space and time*, Acta Math., **135** (1975), pp.57-128.

[13] V.M.Zakalyukin, *Reconstruction of fronts and caustics depending on a parameter and versality of mappings*, J. of Soviet Math., **27** (1984), pp.2713-2735